CAMBRIDGE TRACTS IN MATHEMATICS

General Editors

B. BOLLOBÁS, W. FULTON, A. KATOK, F. KIRWAN, P. SARNAK, B. SIMON, B. TOTARO

196 The Theory of Hardy's Z-Function

CAMBRIDGE TRACTS IN MATHEMATICS

GENERAL EDITORS

B. BOLLOBÁS, W. FULTON, A. KATOK, F. KIRWAN, P. SARNAK, B. SIMON, B. TOTARO

A complete list of books in the series can be found at www.cambridge.org/mathematics. Recent titles include the following:

- 163. Linear and Projective Representations of Symmetric Groups. By A. KLESHCHEV
- 164. The Covering Property Axiom, CPA. By K. CIESIELSKI and J. PAWLIKOWSKI
- 165. Projective Differential Geometry Old and New. By V. OVSIENKO and S. TABACHNIKOV
- 166. The Lévy Laplacian. By M. N. FELLER
- 167. Poincaré Duality Algebras, Macaulay's Dual Systems, and Steenrod Operations. By D. MEYER and L. SMITH
- 168. The Cube-A Window to Convex and Discrete Geometry. By C. ZONG
- 169. Quantum Stochastic Processes and Noncommutative Geometry. By K. B. SINHA and D. GoswAMI
- 170. Polynomials and Vanishing Cycles. By M. TIBĂR
- 171. Orbifolds and Stringy Topology. By A. ADEM, J. LEIDA, and Y. RUAN
- 172. Rigid Cohomology. By B. LE STUM
- 173. Enumeration of Finite Groups. By S. R. BLACKBURN, P. M. NEUMANN, and G. VENKATARAMAN
- 174. Forcing Idealized. By J. ZAPLETAL
- 175. The Large Sieve and its Applications. By E. KOWALSKI
- 176. The Monster Group and Majorana Involutions. By A. A. IVANOV
- 177. A Higher-Dimensional Sieve Method. By H. G. DIAMOND, H. HALBERSTAM, and W. F. GALWAY
- 178. Analysis in Positive Characteristic. By A. N. KOCHUBEI
- 179. Dynamics of Linear Operators. By F. BAYART and É. MATHERON
- 180. Synthetic Geometry of Manifolds. By A. Kock
- 181. Totally Positive Matrices. By A. PINKUS
- 182. Nonlinear Markov Processes and Kinetic Equations. By V. N. KOLOKOLTSOV
- 183. Period Domains over Finite and *p*-adic Fields. By J.-F. DAT, S. ORLIK, and M. RAPOPORT
- 184. Algebraic Theories. By J. ADÁMEK, J. ROSICKÝ, and E. M. VITALE
- 185. Rigidity in Higher Rank Abelian Group Actions I: Introduction and Cocycle Problem. Ву А. КАТОК and V. NIŢICĂ
- 186. Dimensions, Embeddings, and Attractors. By J. C. ROBINSON
- 187. Convexity: An Analytic Viewpoint. By B. SIMON
- 188. Modern Approaches to the Invariant Subspace Problem. By I. CHALENDAR and J. R. PARTINGTON
- 189. Nonlinear Perron Frobenius Theory. By B. LEMMENS and R. NUSSBAUM
- 190. Jordan Structures in Geometry and Analysis. By C.-H. CHU
- Malliavin Calculus for Lévy Processes and Infinite-Dimensional Brownian Motion. By H. Osswald
- 192. Normal Approximations with Malliavin Calculus. By I. NOURDIN and G. PECCATI
- 193. Distribution Modulo One and Diophantine Approximation. By Y. BUGEAUD
- 194. Mathematics of Two-Dimensional Turbulence. By S. KUKSIN and A. SHIRIKYAN
- 195. A Universal Construction for R-free Groups. By I. CHISWELL and T. MÜLLER
- 196. The Theory of Hardy's Z-Function. By A. Ivić
- 197. Induced Representations of Locally Compact Groups. By E. KANIUTH and K. F. TAYLOR
- 198. Topics in Critical Point Theory. By K. PERERA and M. SCHECHTER
- 199. Combinatorics of Minuscule Representations. By R. M. GREEN
- 200. Singularities of the Minimal Model Program. By J. KOLLÁR



G. H. Hardy, 1877-1947

The Theory of Hardy's Z-Function

ALEKSANDAR IVIĆ Univerzitet u Beogradu, Serbia



CAMBRIDGE UNIVERSITY PRESS Cambridge, New York, Melbourne, Madrid, Cape Town, Singapore, São Paulo, Delhi, Mexico City

Cambridge University Press The Edinburgh Building, Cambridge CB2 8RU, UK

Published in the United States of America by Cambridge University Press, New York

www.cambridge.org Information on this title: www.cambridge.org/9781107028838

© Aleksandar Ivić 2013

Frontispiece: G. H. Hardy. Kindly supplied by Trinity College, Cambridge

This publication is in copyright. Subject to statutory exception and to the provisions of relevant collective licensing agreements, no reproduction of any part may take place without the written permission of Cambridge University Press.

First published 2013

Printed and bound in the United Kingdom by the MPG Books Group

A catalog record for this publication is available from the British Library

Library of Congress Cataloguing in Publication data Ivic, A., 1949– author. The theory of Hardy's Z-function / Aleksandar Ivic. pages cm. – (Cambridge tracts in mathematics ; 196) Includes bibliographical references and index. ISBN 978-1-107-02883-8 (hardback) 1. Number theory. I. Title. QA241.I83 2012 512.7 – dc23 2012024804

ISBN 978-1-107-02883-8 Hardback

Cambridge University Press has no responsibility for the persistence or accuracy of URLs for external or third-party internet websites referred to in this publication, and does not guarantee that any content on such websites is, or will remain, accurate or appropriate.

Contents

	Preface		<i>page</i> xi
	Note	ation	XV
1	Definition of $\zeta(s)$, $Z(t)$ and basic notions		1
	1.1	The basic notions	1
	1.2	The functional equation for $\zeta(s)$	3
	1.3	Properties of Hardy's function	6
	1.4	The distribution of zeta-zeros	8
	Not	es	14
2	The zeros on the critical line		21
	2.1	The infinity of zeros on the critical line	21
	2.2	A lower bound for the mean values	23
	2.3	Lehmer's phenomenon	25
	2.4	Gaps between consecutive zeros on the critical line	28
	Not	es	41
3	The Selberg class of <i>L</i> -functions		49
	3.1	The axioms of Selberg's class	49
	3.2	The analogs of Hardy's and Lindelöf's function for ${\cal S}$	51
	3.3	The degree d_F and the invariants of S	52
	3.4	The zeros of functions in \mathcal{S}	56
	Not	es	57
4	The approximate functional equations for $\zeta^k(s)$		61
	4.1	A simple AFE for $\zeta(s)$	61
	4.2	The Riemann-Siegel formula	63
	4.3	The AFE for the powers of $\zeta(s)$	70

viii Contents		
	4.4 The reflection principle4.5 The AFEs with smooth weightsNotes	84 87 94
5	The derivatives of $Z(t)$ 5.1 The θ and Γ functions 5.2 The formula for the derivatives Notes	99 99 101 106
6	 Gram points 6.1 Definition and order of Gram points 6.2 Gram's law 6.3 A mean value result Notes 	109 109 112 115 120
7	The moments of Hardy's function7.1 The asymptotic formula for the moments7.2 RemarksNotes	123 123 130 132
8	The primitive of Hardy's function8.1Introduction8.2The Laplace transform of Hardy's function8.3Proof of Theorem 8.28.4Proof of Theorem 8.3Notes	135 135 138 142 150 153
9	The Mellin transforms of powers of $Z(t)$ 9.1Introduction9.2Some properties of the modified Mellin transforms9.3Analytic continuation of $\mathcal{M}_k(s)$ Notes	157 157 159 164 172
10	Further results on $\mathcal{M}_k(s)$ and $\mathcal{Z}_k(s)$ 10.1 Some relations for $\mathcal{M}_k(s)$ 10.2 Mean square identities for $\mathcal{M}_k(s)$ 10.3 Estimates for $\mathcal{M}_k(s)$ 10.4 Natural boundaries Notes	176 176 180 186 198 201
11	On some problems involving Hardy's function and zeta-moments 11.1 The distribution of values of Hardy's function	206 206

Cambridge University Press
978-1-107-02883-8 - The Theory of Hardy's Z-Function
Aleksandar Ivić
Frontmatter
More information

Contents	ix
11.2 The order of the primitive of Hardy's function	207
11.3 The cubic moment of Hardy's function	209
11.4 Further problems on the distribution of values	213
Notes	219
References	225
Author index	239
Subject index	

Preface

This text has grown out of a mini-course held at the Arctic Number Theory School, University of Helsinki, May 18-25, 2011. The central topic is Hardy's function Z(t), of great importance in the theory of the Riemann zeta-function $\zeta(s)$. It is named after Godfrey Harold ("G. H.") Hardy FRS (1877-1947), who was a prominent English mathematician, well-known for his achievements in number theory and mathematical analysis. Sometimes by Hardy function(s) one denotes the element(s) of Hardy spaces H^p , which are certain spaces of holomorphic functions on the unit disk or the upper half-plane. In this text, however, Hardy's function Z(t) will always denote the function defined by (0) below. It was chosen as the object of study because of its significance in the theory of $\zeta(s)$ and because, initially, considerable material could be presented on the blackboard within the framework of six lectures. Some results, like Theorem 6.7 and the bounds in (4.25) and (4.26) are new, improving on older ones. It is "Hardy's function" which is the thread that holds this work together. I have thought it is appropriate for a monograph because the topic is not as vast as the topic of the Riemann zeta-function itself. Moreover, specialized monographs such as [Iv4], [Lau5], [Mot5] and [Ram] cover in detail other parts of zeta-function theory, but not Z(t). It appears that one volume cannot suffice today to cover all the relevant material concerning $\zeta(s)$, although some of the most important problems (such as the distribution of zeros, in particular the notorious Riemann hypothesis that all complex zeros of $\zeta(s)$ have real parts 1/2), are not settled yet. On the other hand, there seems now to be enough material on Z(t) to warrant the writing of a monograph dedicated solely to it, especially since there is at present a lot of research concerning Z(t) going on.

In the desire to keep the size of the text compact, I had to avoid going into detail in subjects which are connected with zeta-function theory, but are already adequately covered in the literature. These include spectral theory,

Cambridge University Press 978-1-107-02883-8 - The Theory of Hardy's Z-Function Aleksandar Ivić Frontmatter <u>More information</u>

xii

Preface

random matrix theory and representation theory, among others. Proofs of some well-known results such as the first and second derivative tests for exponential integrals (Lemma 2.2 and Lemma 2.3) are omitted, while the proof of, for example, Lemma 7.2 would lead us too much astray. I have tried to include most of the material of the ongoing research on Z(t), thus there is much material from papers which are just published or in print. Starting with Chapter 2, the theory of Hardy's function is systematically developed, hence the title *The Theory of Hardy's Z-Function* seems appropriate. In developing the theory, some modern tools of analytic number theory are systematically used and discussed, such as approximate functional equations, exponential sums and integrals, and integral transforms, to name just a few.

The text is intended for ambitious graduate students, Ph.D. students, and researchers in the field. The prerequisites are basically standard courses in real and complex analysis, but it will help if the reader is familiar with a general text on the Riemann zeta-function, such as the monographs [Tit3], [KaVo] and [Iv1], although efforts have been made to keep the text self-contained. Besides mathematicians, the book will hopefully appeal to physicists whose work is linked to the applications of $\zeta(s)$ in Physics.

The text offers sufficient material for a course of one semester. Except for the introductory Chapter 1, other chapters are basically independent, and various shorter courses may be given based on disposable time and the interest of students.

The material is organized as follows. In Chapter 1 we present the Riemann zeta-function and discuss some of its basic properties. Hardy's function

$$Z(t) := \zeta(\frac{1}{2} + it) \left(\chi(\frac{1}{2} + it) \right)^{-1/2}, \quad \zeta(s) = \chi(s)\zeta(1-s), \tag{0}$$

is introduced and some of its properties are presented. One of the most important aspects of Z(t) is that it is a real-valued function of the real valuable t, and its zeros exactly correspond to the zeros of $\zeta(s)$ on the so-called "critical line" $\mathbb{R}e s = 1/2$. Thus it is very convenient to work with, being real-valued, both theoretically and in conjunction with the calculation of zeta-zeros. It is in Chapter 2 that we discuss the zeros of the zeta-function on the critical line. In particular, we present a variant of Hardy's original proof that there are infinitely many zeta-zeros on $\mathbb{R}e s = 1/2$. Besides that, we discuss Lehmer's phenomenon, namely that the Riemann hypothesis fails if Z(t) has a positive local minimum or a negative local maximum (except at t = 2.47575...) and the unconditional estimation of gaps between the consecutive zeros of Z(t).

Chapter 3 is devoted to the Selberg class S of *L*-functions, of which $\zeta(s)$ can be thought of as a prototype, since the elements of S inherit the intrinsic properties of $\zeta(s)$, namely the Euler product (see (1.1)) and the functional equation

Preface

(see (1.5)). Basic notions and properties of functions in S are presented. The class S (introduced by A. Selberg in 1989) is a natural generalization of $\zeta(s)$, and the modern theory of *L*-functions often focuses on families and classes of *L*-functions containing $\zeta(s)$, such as S.

Chapter 4 is dedicated to AFEs (approximate functional equations) for $\zeta^k(s)$ when $k \in \mathbb{N}$. The AFEs are an essential tool in the theory of *L*-functions, approximating a Dirichlet series by a finite number of Dirichlet polynomials. For the AFEs that we treat, the accent is on the methods of proof, of which there are several. From an AFE for $\zeta^k(s)$ one then deduces an AFE for $Z^k(t)$.

In Chapter 5 we provide formulas for the derivatives $Z^{(k)}(t)$, and in Chapter 6 we deal with Gram points g_n which satisfy $\theta(g_n) = \pi n$, where $Z(t) = e^{i\theta(t)}\zeta(\frac{1}{2} + it)$ and $\theta(t)$ is given by (1.19). We also discuss the so-called "Gram's law" related to Gram points.

In Chapter 7 we give formulas for the *k*th moment of Z(t), that is, for the integral $\int_0^T Z^k(t) dt$, where $k \in \mathbb{N}$. This is achieved by employing an AFE with smooth weights.

M. Jutila's recent formula for the oscillating function $F(T) = \int_0^T Z(t) dt$ is given in detail in Chapter 8. It furnishes the results $F(T) = O(T^{1/4})$ and $F(T) = \Omega_{\pm}(T^{1/4})$, thereby determining the true order of magnitude of F(T). This approach is different from the author's work and from the recent work of M. Korolev [Kor3], [Kor4], who also obtained an explicit formula for F(T).

The Mellin transforms $(k \in \mathbb{N})$

$$\mathcal{Z}_k(s) = \int_1^\infty |\zeta(\frac{1}{2} + ix)|^{2k} x^{-s} \, \mathrm{d}x, \quad M_k(s) = \int_1^\infty Z^k(x) x^{-s} \, \mathrm{d}x$$

form the subject of Chapter 9 and Chapter 10. In general, classical integral transforms, in particular Fourier, Laplace and Mellin transforms, play an important rôle in analytic number theory. Analytic continuation, pointwise and mean square estimates of $Z_k(s)$, $M_k(s)$ are given in detail, by using various techniques of analytic number theory. The importance of these functions is their connections with power moments of $|\zeta(\frac{1}{2} + it)|$, one of the most important topics of zeta-function theory.

The concluding Chapter 11 focuses on some problems involving Hardy's function and zeta moments. I hope that this will be of interest to all who want to do research in areas connected to these topics. This is in the spirit of P. Erdős's motto: "Prove and conjecture!".

Each chapter is followed by "Notes", where many references, comments and remarks are given. This seemed preferable to burdening the body of the text with footnotes, parentheses, etc. On many occasions dates of birth and death are given for the mathematicians mentioned in the text.

xiii

xiv

Preface

A full bibliography of all the relevant works mentioned in the text is to be found at the end of the text.

Acknowledgements. Several experts in the field have read (parts of) the text, and made valuable remarks. I wish to thank all of them. In alphabetical order they are: H. Bui, J. Kaczorowski, M. Korolev, Y. Motohashi, A. Perelli, I. Rezvyakova, T. Trudgian and N. Watt. Special thanks go to M. Jutila.

I also wish to thank the staff of Cambridge University Press for their unfailing courtesy and help.

Notation

Owing to the nature of this text, absolute consistency in notation could not be attained, although whenever possible standard notation is used. Notation used commonly through the text is explained there, while specific notation introduced in the proof of a theorem or lemma is given at the proper place in the body of the text.

k, l, m, n, \ldots	Natural numbers (positive integers).
p	A generic prime number.
$\mathbb{N}, \mathbb{Z}, \mathbb{R}, \mathbb{C}$	The sets of natural numbers, integers, real and com-
	plex numbers, respectively.
A, B, C, C_1, \ldots	Absolute, positive constants (not necessarily the
	same ones at each occurrence).
ε	An arbitrarily small positive number, not necessarily
	the same one at each occurrence.
s, z, w	Complex variables ($\mathbb{R}e s$ and $\mathbb{I}m s$ denote the real and
	imaginary part of s, respectively; common notation
	is $\sigma = \mathbb{R}e s$ and $t = \mathbb{I}m s$).
<i>t</i> , <i>x</i> , <i>y</i>	Real variables.
$\operatorname{Res}_{s=s_0} F(s)$	Denotes the residue of $F(s)$ at the point $s = s_0$.
$\zeta(s)$	The Riemann zeta-function is defined by $\zeta(s) =$
	$\sum_{n=1}^{\infty} n^{-s}$ for $\mathbb{R}es > 1$ and otherwise by analytic
	continuation.
$\Gamma(s)$	$=\int_0^\infty x^{s-1} e^{-x} dx$ for $\mathbb{R}e s > 0$, otherwise by ana-
	lytic continuation by $s\Gamma(s) = \Gamma(s+1)$. This is the
	Euler gamma-function.
γ	Euler's constant $\gamma = -\Gamma'(1) = 0.5772157\ldots$

xvi	Notation
$\chi(s)$	The function defined by $\zeta(s) = \chi(s)\zeta(1-s)$, so that by the functional equation for $\zeta(s)$ we have $\chi(s) = (2\pi)^s/(2\Gamma(s)\cos(\pi s/2))$.
Z(t)	Hardy's function, $Z(t) = \zeta(\frac{1}{2} + it)(\chi(\frac{1}{2} + it))^{-1/2}$.
$\theta(t)$	For real t defined as $\theta(t) = \mathbb{I}m \left\{ \log \Gamma(\frac{1}{4} + \frac{1}{2}it) \right\} - \frac{1}{2}t \log \pi$.
$\rho = \beta + i\gamma$ $N(T)$	A complex zero of $\zeta(s)$; $\beta = \mathbb{R}e \rho$, $\gamma = \mathbb{I}m \rho$. The number of zeros $\rho = \beta + i\gamma$ of $\zeta(s)$, counted with multiplicities for which $0 < \gamma < T$
S(T)	$= \frac{1}{2} \arg \zeta(\frac{1}{2} + iT).$
$\mu(\sigma)$	For real σ defined as $\mu(\sigma) = \limsup_{t \to \infty} \frac{\log \zeta(\sigma + it) }{\log t}$.
$\exp(z)$	$= e^{z}$.
e(z)	$=e^{2\pi i z}$.
$\log x$	$= \text{Log}_{e} x \equiv \ln x.$
[<i>x</i>]	The greatest integer not exceeding the real number x.
$\{x\}$	= x - [x], the fractional part of x.
$\sum_{n \le x} f(n)$	A sum taken over all natural numbers <i>n</i> not exceeding
	<i>x</i> ; the empty sum is defined to be equal to zero.
\prod_{i}	The product taken over all possible values of the
5	index j ; the empty product is defined to be unity.
$d_k(n)$	The number of ways <i>n</i> can be written as a product of $k \ge 2$ fixed factors; $d_2(n) = d(n)$ is the number of
	divisors of <i>n</i> .
$\Delta(x)$	= $\sum_{n \le x} d(n) - x(\log x + 2\gamma - 1)$, the error term in the Dirichlet divisor problem.
$f(x) \sim g(x)$	Means $\lim_{x\to x_0} f(x)/g(x) = 1$, as $x \to x_0$, with x_0 not necessarily finite.
f(x) = O(g(x))	Means $ f(x) \le Cg(x)$ for $x \ge x_0$ and some constant
	C > 0. Here $f(x)$ is a complex function of the real
	variable <i>x</i> and $g(x)$ is a positive function of <i>x</i> for $x \ge x$
	$x_0. f(x) = O_{\alpha,\beta,\dots}(g(x))$ means the implied constant
	depends on α , β ,
$f(x) \ll g(x)$	Means the same as $f(x) = O(g(x))$. Likewise
	$f(x) \ll_{\alpha,\beta,\dots} g(x)$ means the implied constant depends on α, β, \dots .
$f(x) \gg g(x)$	Means the same as $g(x) = O(f(x))$.
$f(x) \asymp g(x)$	Means that both $f(x) \ll g(x)$ and $g(x) \ll f(x)$
	hold.

Notation

xvii

(a,b)	Means the interval $a < x < b$.
[a,b]	Means the interval $a \le x \le b$.
$C^{r}[a,b]$	The class of functions having a continuous <i>r</i> th
	derivative in $[a, b]$.
$L^p(a,b)$	The class of measurable functions $f(x)$ such that
	$\int_{a}^{b} f(x) ^{p} dx$ is finite.
f(x) = o(g(x)) as	Means $\lim_{x\to x_0} f(x)/g(x) = 0$, with x_0 possibly
$x \to x_0$	infinite.
$f(x) = \Omega(g(x))$	Means that $f(x) = o(g(x))$ does not hold when
	$x \to \infty$.
$f(x) = \Omega_+(g(x))$	Means that there exists a suitable constant $C > 0$
	such that $f(x) > Cg(x)$ holds for some arbitrarily
	large values of x.
$f(x) = \Omega_{-}(g(x))$	Means that there exists a suitable constant $C > 0$
	such that $f(x) < -Cg(x)$ holds for some arbitrarily
	large values of x.
$f(x) = \Omega_{\pm}(g(x))$	Means that both $f(x) = \Omega_+(g(x))$ and
	$f(x) = \Omega_{-}(g(x))$ hold.
$\int C(z) dz$	$\frac{c+iT}{c+iT}$
$J_{(c)} G(s) ds$	$= \lim_{T \to \infty} \int_{c-iT} G(s) \mathrm{d}s.$
	$\int_{0}^{\infty} \mathbf{Z} k(x) x^{-\delta} dx \qquad (h \in \mathbb{N})$
$\mathcal{M}_k(s)$	$= \int Z^{n}(x)x^{n-1} dx (k \in \mathbb{N}).$
$\mathcal{Z}(\mathbf{s})$	$= \int z ^2 (1 + iz) ^{2k} z^{-s} dz (k \in \mathbb{N})$
$\mathcal{L}_k(s)$	$= \int_{1} \zeta(\frac{1}{2} + ix) x dx (k \in \mathbb{N}).$
AFE	Means "approximate functional equation".