

Chapter 2

Duality and a Farkas lemma for integer programs

Jean B. Lasserre

Abstract We consider the integer program $\max\{c'x \mid Ax = b, x \in \mathbf{N}^n\}$. A formal parallel between linear programming and continuous integration, and discrete summation, shows that a natural duality for integer programs can be derived from the \mathbf{Z} -transform and Brion and Vergne's counting formula. Along the same lines, we also provide a discrete Farkas lemma and show that the existence of a nonnegative integral solution $x \in \mathbf{N}^n$ to $Ax = b$ can be tested via a linear program.

Key words: Integer programming, counting problems, duality

2.1 Introduction


In this paper we are interested in a comparison between linear and integer programming, and particularly in a *duality* perspective. So far, and to the best of our knowledge, the duality results available for integer programs are obtained via the use of *subadditive* functions as in Wolsey [21], for example, and the smaller class of *Chvátal* and *Gomory* functions as in Blair and Jeroslow [6], for example (see also Schrijver [19, pp. 346–353]). For more details the interested reader is referred to [1, 6, 19, 21] and the many references therein. However, as subadditive, Chvátal and Gomory functions are only defined implicitly from their properties, the resulting dual problems defined in [6] or [21] are conceptual in nature and Gomory functions are used to generate valid inequalities for the primal problem.

We claim that another natural *duality* for integer programs can be derived from the \mathbf{Z} -transform (or generating function) associated with the *counting*

Jean B. Lasserre
LAAS-CNRS, 7 Avenue du Colonel Roche, 31077 Toulouse Cédex 4, FRANCE
e-mail: lasserre@laas.fr

version (defined below) of the integer program. Results for counting problems, notably by Barvinok [4], Barvinok and Pommersheim [5], Khovanskii and Pukhlikov [12], and in particular, Brion and Vergne’s counting formula [7], will prove especially useful.

For this purpose, we will consider the four related problems \mathbf{P} , \mathbf{P}_d , \mathbf{I} and \mathbf{I}_d displayed in the diagram below, in which the integer program \mathbf{P}_d appears in the upper right corner.



Problem \mathbf{I} (in which ds denotes the Lebesgue measure on the affine variety $\{x \in \mathbf{R}^n \mid Ax = b\}$ that contains the convex polyhedron $\Omega(b)$) is the *integration* version of the linear program \mathbf{P} , whereas Problem \mathbf{I}_d is the *counting* version of the (discrete) integer program \mathbf{P}_d .

Why should these four problems help in analyzing \mathbf{P}_d ? Because first, \mathbf{P} and \mathbf{I} , as well as \mathbf{P}_d and \mathbf{I}_d , are simply related, and in the same manner. Next, as we will see, the nice and complete duality results available for \mathbf{P} , \mathbf{I} and \mathbf{I}_d extend in a natural way to \mathbf{P}_d .

2.1.1 Preliminaries

In fact, \mathbf{I} and \mathbf{I}_d are the respective formal analogues in the algebra $(+, \times)$ of \mathbf{P} and \mathbf{P}_d in the algebra (\oplus, \times) , where in the latter, the addition $a \oplus b$ stands for $\max(a, b)$; indeed, the “max” in \mathbf{P} and \mathbf{P}_d can be seen as an *idempotent* integral (or *Maslov integral*) in this algebra (see, for example, Litvinov *et al.* [17]). For a nice parallel between results in probability $((+, \times)$ algebra) and optimization $((\max, +)$ algebra), the reader is referred to Bacelli *et al.* [3, Section 9].

Moreover, \mathbf{P} and \mathbf{I} , as well as \mathbf{P}_d and \mathbf{I}_d , are simply related via

$$\varepsilon^{f(b,c)} = \lim_{r \rightarrow \infty} \hat{f}(b, rc)^{1/r}; \quad \varepsilon^{f_d(b,c)} = \lim_{r \rightarrow \infty} \hat{f}_d(b, rc)^{1/r}. \quad (2.1)$$

Equivalently, by continuity of the logarithm,

$$f(b, c) = \lim_{r \rightarrow \infty} \frac{1}{r} \ln \widehat{f}(b, rc); \quad f_d(b, c) = \lim_{r \rightarrow \infty} \frac{1}{r} \ln \widehat{f}_d(b, rc), \quad (2.2)$$

a relationship that will be useful later.

Next, concerning *duality*, the standard *Legendre-Fenchel* transform which yields the usual dual LP of \mathbf{P} ,

$$\mathbf{P}^* \rightarrow \min_{\lambda \in \mathbf{R}^m} \{b' \lambda \mid A' \lambda \geq c\}, \quad (2.3)$$

has a natural analogue for integration, the *Laplace transform*, and thus the *inverse Laplace transform* problem (that we call \mathbf{I}^*) is the formal analogue of \mathbf{P}^* and provides a nice duality for integration (although not usually presented in these terms). Finally, the \mathbf{Z} -transform is the obvious analogue for *summation* of the Laplace transform for integration. We will see that in the light of recent results in *counting problems*, it is possible to establish a nice duality for \mathbf{I}_d in the same vein as the duality for (continuous) integration and by (2.2), it also provides a powerful tool for analyzing the integer program \mathbf{P}_d .

2.1.2 Summary of content

(a) We first review the duality principles that are available for \mathbf{P} , \mathbf{I} and \mathbf{I}_d and underline the parallels and connections between them. In particular, a fundamental difference between the continuous and discrete cases is that in the former, the data appear as *coefficients* of the dual variables whereas in the latter, the *same* data appear as *exponents* of the dual variables. Consequently, the (discrete) \mathbf{Z} -transform has many more *poles* than the Laplace transform. Whereas the Laplace transform has only *real* poles, the \mathbf{Z} -transform has additional *complex* poles associated with each real pole, which induces some *periodic* behavior, a well-known phenomenon in number theory where the \mathbf{Z} -transform (or *generating function*) is a standard tool (see, for example, Iosevich [11], Mitrinović *et al.* [18]). So, if the procedure of inverting the Laplace transform or the \mathbf{Z} -transform (that is, solving the dual problems \mathbf{I}^* and \mathbf{I}_d^*) is basically of the same nature, that is, a complex integral, it is significantly more complicated in the discrete case, due to the presence of these additional complex poles.

(b) Then we use results from (a) to analyze the discrete optimization problem \mathbf{P}_d . Central to the analysis is Brion and Vergne's inverse formula [7] for counting problems. In particular, we provide a closed-form expression for the optimal value $f_d(b, c)$ which highlights the special role played by the so-called *reduced costs* of the linear program \mathbf{P} and the *complex poles* of the \mathbf{Z} -transform associated with each basis of the linear program \mathbf{P} . We also show that each basis B of the linear program \mathbf{P} provides exactly $\det(B)$

complex *dual* vectors in \mathbf{C}^m , the complex (periodic) analogues for \mathbf{P}_d of the unique dual vector in \mathbf{R}^m for \mathbf{P} , associated with the basis B . As in linear programming (but in a more complicated way), the optimal value $f_d(b, c)$ of \mathbf{P}_d can be found by inspection of (certain sums of) reduced costs associated with each vertex of $\Omega(b)$.

(c) We also provide a *discrete* Farkas lemma for the existence of nonnegative integral solutions $x \in \mathbf{N}^n$ to $Ax = b$. Its form also confirms the special role of the \mathbf{Z} -transform described earlier. Moreover, it allows us to check the existence of a nonnegative integral solution by solving a related linear program.

2.2 Duality for the continuous problems \mathbf{P} and \mathbf{I}

With $A \in \mathbf{R}^{m \times n}$ and $b \in \mathbf{R}^m$, let $\Omega(b)\mathbf{R}^n$ be the convex polyhedron

$$\Omega(b) := \{x \in \mathbf{R}^n \mid Ax = b; \quad x \geq 0\}, \quad (2.4)$$

and consider the standard linear program (LP)

$$\mathbf{P}: \quad f(b, c) := \max\{c'x \mid Ax = b; \quad x \geq 0\} \quad (2.5)$$

with $c \in \mathbf{R}^n$, and its associated *integration* version

$$\mathbf{I}: \quad \hat{f}(b, c) := \int_{\Omega(b)} \varepsilon^{c'x} ds \quad (2.6)$$

where ds is the Lebesgue measure on the affine variety $\{x \in \mathbf{R}^n \mid Ax = b\}$ that contains the convex polyhedron $\Omega(b)$.

For a vector c and a matrix A we denote by c' and A' their respective transposes. We also use both the notation $c'x$ and $\langle c, x \rangle$ for the usual scalar product of two vectors c and x . We assume that both $A \in \mathbf{R}^{m \times n}$ and $b \in \mathbf{R}^m$ have rational entries.

2.2.1 Duality for \mathbf{P}

It is well known that the standard duality for (2.5) is obtained from the *Legendre-Fenchel* transform $F(\cdot, c) : \mathbf{R}^m \rightarrow \mathbf{R}$ of the value function $f(b, c)$ with respect to b , that is, here (as $y \mapsto f(y, c)$ is concave)

$$\lambda \mapsto F(\lambda, c) := \inf_{y \in \mathbf{R}^m} \langle \lambda, y \rangle - f(y, c), \quad (2.7)$$

which yields the usual dual LP problem

$$\mathbf{P}^* \rightarrow \inf_{\lambda \in \mathbf{R}^m} \langle \lambda, b \rangle - F(\lambda, c) = \min_{\lambda \in \mathbf{R}^m} \{b'\lambda \mid A'\lambda \geq c\}. \quad (2.8)$$

2.2.2 Duality for integration

Similarly, the analogue for integration of the Fenchel transform is the two-sided Laplace transform $\widehat{F}(\cdot, c) : \mathbf{C}^m \rightarrow \mathbf{C}$ of $\widehat{f}(b, c)$, given by

$$\lambda \mapsto \widehat{F}(\lambda, c) := \int_{\mathbf{R}^m} \varepsilon^{-\langle \lambda, y \rangle} \widehat{f}(y, c) dy. \quad (2.9)$$

It turns out that developing (2.9) yields

$$\widehat{F}(\lambda, c) = \prod_{k=1}^n \frac{1}{(A'\lambda - c)_k} \quad \text{whenever } \mathbf{Re}(A'\lambda - c) > 0, \quad (2.10)$$

(see for example [7, p. 798] or [13]). Thus $\widehat{F}(\lambda, c)$ is well-defined provided

$$\mathbf{Re}(A'\lambda - c) > 0, \quad (2.11)$$

and $\widehat{f}(b, c)$ can be computed by solving the *inverse* Laplace transform problem, which we call the (integration) *dual* problem \mathbf{I}^* of (2.12), that is,

$$\begin{aligned} \mathbf{I}^* \rightarrow \widehat{f}(b, c) &:= \frac{1}{(2i\pi)^m} \int_{\gamma-i\infty}^{\gamma+i\infty} \varepsilon^{\langle b, \lambda \rangle} \widehat{F}(\lambda, c) d\lambda \\ &= \frac{1}{(2i\pi)^m} \int_{\gamma-i\infty}^{\gamma+i\infty} \frac{\varepsilon^{\langle b, \lambda \rangle}}{\prod_{k=1}^n (A'\lambda - c)_k} d\lambda, \end{aligned} \quad (2.12)$$

where $\gamma \in \mathbf{R}^m$ is fixed and satisfies $A'\gamma - c > 0$. Incidentally, observe that the domain of definition (2.11) of $\widehat{F}(\cdot, c)$ is precisely the interior of the feasible set of the dual problem \mathbf{P}^* in (2.8). We will comment more on this and the link with the logarithmic barrier function for linear programming (see Section 2.2.5 below).

We may indeed call \mathbf{I}^* a dual problem of \mathbf{I} as it is defined on the space \mathbf{C}^m of variables $\{\lambda_k\}$ associated with the nontrivial constraints $Ax = b$; notice that we also retrieve the standard “ingredients” of the dual optimization problem \mathbf{P}^* , namely $b'\lambda$ and $A'\lambda - c$.

2.2.3 Comparing \mathbf{P}, \mathbf{P}^* and \mathbf{I}, \mathbf{I}^*

One may compute $\widehat{f}(b, c)$ directly using Cauchy residue techniques. That is, one may compute the integral (2.12) by successive one-dimensional complex integrals with respect to one variable λ_k at a time (for example starting with $\lambda_1, \lambda_2, \dots$) and by repeated application of Cauchy’s Residue Theorem

[8]. This is possible because the integrand is a rational fraction, and after application of Cauchy's Residue Theorem at step k with respect to λ_k , the output is still a rational fraction of the remaining variables $\lambda_{k+1}, \dots, \lambda_m$. For more details the reader is referred to Lasserre and Zeron [13]. It is not difficult to see that the whole procedure is a summation of partial results, each of them corresponding to a (multi-pole) vector $\hat{\lambda} \in \mathbf{R}^m$ that annihilates m terms of n products in the denominator of the integrand.

This is formalized in the nice formula of Brion and Vergne [7, Proposition 3.3 p. 820] that we describe below. For the interested reader, there are several other nice closed-form formulae for $\hat{f}(b, c)$, notably by Barvinok [4], Barvinok and Pommersheim [5], and Khovanskii and Pukhlikov [12].

2.2.4 The continuous Brion and Vergne formula

The material in this section is taken from [7]. To explain the closed-form formula of Brion and Vergne we need some notation.

Write the matrix $A \in \mathbf{R}^{m \times n}$ as $A = [A_1 | \dots | A_n]$ where $A_j \in \mathbf{R}^m$ denotes the j -th column of A for all $j = 1, \dots, n$. With $\Delta := (A_1, \dots, A_n)$ let $C(\Delta) \subset \mathbf{R}^m$ be the closed convex cone generated by Δ . Let $\Lambda \subseteq \mathbf{Z}^m$ be a lattice.

A subset σ of $\{1, \dots, n\}$ is called a *basis* of Δ if the sequence $\{A_j\}_{j \in \sigma}$ is a basis of \mathbf{R}^m , and the set of bases of Δ is denoted by $\mathcal{B}(\Delta)$. For $\sigma \in \mathcal{B}(\Delta)$ let $C(\sigma)$ be the cone generated by $\{A_j\}_{j \in \sigma}$. With any $y \in C(\Delta)$ associate the intersection of all cones $C(\sigma)$ which contain y . This defines a subdivision of $C(\Delta)$ into polyhedral cones. The interiors of the maximal cones in this subdivision are called *chambers* in Brion and Vergne [7]. For every $y \in \gamma$, the convex polyhedron $\Omega(y)$ in (2.4) is *simple*. Next, for a chamber γ (whose closure is denoted by $\bar{\gamma}$), let $\mathcal{B}(\Delta, \gamma)$ be the set of bases σ such that γ is contained in $C(\sigma)$, and let $\mu(\sigma)$ denote the volume of the convex polytope $\{\sum_{j \in \sigma} t_j A_j \mid 0 \leq t_j \leq 1\}$ (normalized so that $\text{vol}(\mathbf{R}^m/\Lambda) = 1$). Observe that for $b \in \bar{\gamma}$ and $\sigma \in \mathcal{B}(\Delta, \gamma)$ we have $b = \sum_{j \in \sigma} x_j(\sigma) A_j$ for some $x_j(\sigma) \geq 0$. Therefore the vector $x(\sigma) \in \mathbf{R}_+^n$, with $x_j(\sigma) = 0$ whenever $j \notin \sigma$, is a *vertex* of the polytope $\Omega(b)$. In linear programming terminology, the bases $\sigma \in \mathcal{B}(\Delta, \gamma)$ correspond to the *feasible bases* of the linear program \mathbf{P} . Denote by V the subspace $\{x \in \mathbf{R}^n \mid Ax = 0\}$. Finally, given $\sigma \in \mathcal{B}(\Delta)$, let $\pi^\sigma \in \mathbf{R}^m$ be the row vector that solves $\pi^\sigma A_j = c_j$ for all $j \in \sigma$. A vector $c \in \mathbf{R}^n$ is said to be *regular* if $c_j - \pi^\sigma A_j \neq 0$ for all $\sigma \in \mathcal{B}(\Delta)$ and all $j \notin \sigma$. Let $c \in \mathbf{R}^n$ be regular with $-c$ in the interior of the dual cone $(\mathbf{R}_+^n \cap V)^*$ (which is the case if $A'u > c$ for some $u \in \mathbf{R}^m$). Then, with $\Lambda = \mathbf{Z}^m$, Brion and Vergne's formula [7, Proposition 3.3, p. 820] states that

$$\hat{f}(b, c) = \sum_{\sigma \in \mathcal{B}(\Delta, \gamma)} \frac{\varepsilon^{(c, x(\sigma))}}{\mu(\sigma) \prod_{k \notin \sigma} (-c_k + \pi^\sigma A_k)} \quad \forall b \in \bar{\gamma}. \quad (2.13)$$

Notice that in linear programming terminology, $c_k - \pi^\sigma A_k$ is simply the so-called *reduced cost* of the variable x_k , with respect to the basis $\{A_j\}_{j \in \sigma}$. Equivalently, we can rewrite (2.13) as

$$\widehat{f}(b, c) = \sum_{x(\sigma): \text{vertex of } \Omega(b)} \frac{\varepsilon^{\langle c, x(\sigma) \rangle}}{\mu(\sigma) \prod_{k \notin \sigma} (-c_k + \pi^\sigma A_k)}. \quad (2.14)$$

Thus $\widehat{f}(b, c)$ is a weighted *summation* over the vertices of $\Omega(b)$ whereas $f(b, c)$ is a *maximization* over the vertices (or a summation with $\oplus \equiv \max$).

So, if c is replaced by rc and $x(\sigma^*)$ denotes the vertex of $\Omega(b)$ at which $c'x$ is maximized, we obtain

$$\widehat{f}(b, rc)^{1/r} = \varepsilon^{\langle c, x(\sigma^*) \rangle} \left[\sum_{x(\sigma): \text{vertex of } \Omega(b)} \frac{\varepsilon^{r \langle c, x(\sigma) - x(\sigma^*) \rangle}}{r^{n-m} \mu(\sigma) \prod_{k \notin \sigma} (-c_k + \pi^\sigma A_k)} \right]^{\frac{1}{r}}$$

from which it easily follows that

$$\lim_{r \rightarrow \infty} \ln \widehat{f}(b, rc)^{1/r} = \langle c, x(\sigma^*) \rangle = \max_{x \in \Omega(b)} \langle c, x \rangle = f(b, c),$$

as indicated in (2.2).

2.2.5 The logarithmic barrier function

It is also worth noticing that

$$\begin{aligned} \widehat{f}(b, rc) &= \frac{1}{(2i\pi)^m} \int_{\gamma_r - i\infty}^{\gamma_r + i\infty} \frac{\varepsilon^{\langle b, \lambda \rangle}}{\prod_{k=1}^n (A' \lambda - rc)_k} d\lambda \\ &= \frac{1}{(2i\pi)^m} \int_{\gamma - i\infty}^{\gamma + i\infty} \frac{r^{m-n} \varepsilon^{\langle rb, \lambda \rangle}}{\prod_{k=1}^n (A' \lambda - c)_k} d\lambda \end{aligned}$$

with $\gamma_r = r\gamma$ and we can see that (up to the constant $(m-n) \ln r$) the logarithm of the integrand is simply the well-known *logarithmic barrier function*

$$\lambda \mapsto \phi_\mu(\lambda, b) = \mu^{-1} \langle b, \lambda \rangle - \sum_{j=1}^n \ln (A' \lambda - c)_j,$$

with parameter $\mu := 1/r$, of the dual problem \mathbf{P}^* (see for example den Hertog [9]). This should not come as a surprise as a self-concordant barrier function $\phi_K(x)$ of a cone $K \subset \mathbf{R}^n$ is given by the logarithm of the Laplace transform $\int_{K^*} \varepsilon^{-\langle x, s \rangle} ds$ of its dual cone K^* (see for example Güler [10], Truong and Tunçel [20]).

Thus, when $r \rightarrow \infty$, minimizing the exponential logarithmic barrier function on its domain in \mathbf{R}^m yields the same result as taking its residues.

2.2.6 Summary

The parallel between \mathbf{P}, \mathbf{P}^* and \mathbf{I}, \mathbf{I}^* is summarized below.

Fenchel-duality

$$f(b, c) := \max_{Ax=b; x \geq 0} c'x$$

$$F(\lambda, c) := \inf_{y \in \mathbf{R}^m} \{\lambda'y - f(y, c)\}$$

with : $A'\lambda - c \geq 0$

$$f(b, c) = \min_{\lambda \in \mathbf{R}^m} \{\lambda'b - F(\lambda, c)\}$$

$$= \min_{\lambda \in \mathbf{R}^m} \{b'\lambda \mid A'\lambda \geq c\}$$

Simplex algorithm \rightarrow

vertices of $\Omega(b)$

$\rightarrow \max c'x$ over vertices.

Laplace-duality

$$\widehat{f}(b, c) := \int_{Ax=b; x \geq 0} \varepsilon^{c'x} ds$$

$$\begin{aligned} \widehat{F}(\lambda, c) &:= \int_{\mathbf{R}^m} \varepsilon^{-\lambda'y} \widehat{f}(y, c) dy \\ &= \frac{1}{\prod_{k=1}^n (A'\lambda - c)_k} \end{aligned}$$

with : $\mathbf{Re}(A'\lambda - c) > 0$

$$\begin{aligned} \widehat{f}(b, c) &= \frac{1}{(2i\pi)^m} \int_{\Gamma} \varepsilon^{\lambda'b} \widehat{F}(\lambda, c) d\lambda \\ &= \frac{1}{(2i\pi)^m} \int_{\Gamma} \frac{\varepsilon^{\lambda'b}}{\prod_{k=1}^n (A'\lambda - c)_k} d\lambda \end{aligned}$$

Cauchy's Residue \rightarrow

poles of $\widehat{F}(\lambda, c)$

$\rightarrow \sum \varepsilon^{c'x}$ over vertices.

2.3 Duality for the discrete problems \mathbf{I}_d and \mathbf{P}_d

In the respective *discrete* analogues \mathbf{P}_d and \mathbf{I}_d of (2.5) and (2.6) one replaces the positive cone \mathbf{R}_+^n by \mathbf{N}^n (or $\mathbf{R}_+^n \cap \mathbf{Z}^n$), that is, (2.5) becomes the integer program

$$\mathbf{P}_d: f_d(b, c) := \max \{c'x \mid Ax = b; x \in \mathbf{N}^n\} \quad (2.15)$$

whereas (2.6) becomes a summation over $\mathbf{N}^n \cap \Omega(b)$, that is,

$$\mathbf{I}_d: \widehat{f}_d(b, c) := \sum \{ \varepsilon^{c'x} \mid Ax = b; \quad x \in \mathbf{N}^n \}. \quad (2.16)$$

We here assume that $A \in \mathbf{Z}^{m \times n}$ and $b \in \mathbf{Z}^m$, which implies in particular that the lattice $\Lambda := A(\mathbf{Z}^n)$ is a sublattice of \mathbf{Z}^m ($\Lambda \subseteq \mathbf{Z}^m$). Note that b in (2.15) and (2.16) is necessarily in Λ .

In this section we are concerned with what we call the “dual” problem \mathbf{I}_d^* of \mathbf{I}_d , the discrete analogue of the dual \mathbf{I}^* of \mathbf{I} , and its link with the discrete optimization problem \mathbf{P}_d .

2.3.1 The \mathbf{Z} -transform

The natural discrete analogue of the Laplace transform is the so-called \mathbf{Z} -transform. Therefore with $\widehat{f}_d(b, c)$ we associate its (two-sided) \mathbf{Z} -transform $\widehat{F}_d(\cdot, c) : \mathbf{C}^m \rightarrow \mathbf{C}$ defined by

$$z \mapsto \widehat{F}_d(z, c) := \sum_{y \in \mathbf{Z}^m} z^{-y} \widehat{f}_d(y, c), \quad (2.17)$$

where the notation z^y with $y \in \mathbf{Z}^m$ stands for $z_1^{y_1} \cdots z_m^{y_m}$. Applying this definition yields

$$\begin{aligned} \widehat{F}_d(z, c) &= \sum_{y \in \mathbf{Z}^m} z^{-y} \widehat{f}_d(y, c) \\ &= \sum_{y \in \mathbf{Z}^m} z^{-y} \left[\sum_{x \in \mathbf{N}^n; Ax=y} \varepsilon^{c'x} \right] \\ &= \sum_{x \in \mathbf{N}^n} \varepsilon^{c'x} \left[\sum_{y=Ax} z_1^{-y_1} \cdots z_m^{-y_m} \right] \\ &= \sum_{x \in \mathbf{N}^n} \varepsilon^{c'x} z_1^{-(Ax)_1} \cdots z_m^{-(Ax)_m} \\ &= \prod_{k=1}^n \frac{1}{(1 - \varepsilon^{c_k} z_1^{-A_{1k}} z_2^{-A_{2k}} \cdots z_m^{-A_{mk}})} \\ &= \prod_{k=1}^n \frac{1}{(1 - \varepsilon^{c_k} z^{-A_k})}, \end{aligned} \quad (2.18)$$

which is well-defined provided

$$|z_1^{A_{1k}} \cdots z_m^{A_{mk}}| (= |z^{A_k}|) > \varepsilon^{c_k} \quad \forall k = 1, \dots, n. \quad (2.19)$$

Observe that the domain of definition (2.19) of $\widehat{F}_d(\cdot, c)$ is the *exponential* version of (2.11) for $\widehat{F}(\cdot, c)$. Indeed, taking the real part of the logarithm in (2.19) yields (2.11).

2.3.2 The dual problem \mathbf{I}_d^*

Therefore the value $\widehat{f}_d(b, c)$ is obtained by solving the *inverse Z-transform* problem \mathbf{I}_d^* (that we call the *dual* of \mathbf{I}_d)

$$\widehat{f}_d(b, c) = \frac{1}{(2i\pi)^m} \int_{|z_1|=\gamma_1} \cdots \int_{|z_m|=\gamma_m} \widehat{F}_d(z) z^{b-e_m} dz_m \cdots dz_1, \quad (2.20)$$

where e_m is the unit vector of \mathbf{R}^m and $\gamma \in \mathbf{R}^m$ is a (fixed) vector that satisfies $\gamma_1^{A_{1k}} \gamma_2^{A_{2k}} \cdots \gamma_m^{A_{mk}} > \varepsilon^{c_k}$ for all $k = 1, \dots, n$. We may indeed call \mathbf{I}_d^* the *dual problem* of \mathbf{I}_d as it is defined on the space \mathbf{Z}^m of dual variables z_k associated with the nontrivial constraints $Ax = b$ of the primal problem \mathbf{I}_d . We also have the following parallel.

Continuous Laplace-duality

$$\widehat{f}(b, c) := \int_{Ax=b; x \in \mathbf{R}_+^n} \varepsilon^{c'x} ds$$

$$\widehat{F}(\lambda, c) := \int_{\mathbf{R}^m} \varepsilon^{-\lambda'y} \widehat{f}(y, c) dy$$

$$= \prod_{k=1}^n \frac{1}{(A'\lambda - c)_k}$$

with $\mathbf{Re}(A'\lambda - c) > 0$.

Discrete Z-duality

$$\widehat{f}_d(b, c) := \sum_{Ax=b; x \in \mathbf{N}^n} \varepsilon^{c'x}$$

$$\widehat{F}_d(z, c) := \sum_{y \in \mathbf{Z}^m} z^{-y} \widehat{f}_d(y, c)$$

$$= \prod_{k=1}^n \frac{1}{1 - \varepsilon^{c_k} z^{-A_k}}$$

with $|z^{A_k}| > \varepsilon^{c_k}$, $k = 1, \dots, n$.

2.3.3 Comparing \mathbf{I}^* and \mathbf{I}_d^*

Observe that the dual problem \mathbf{I}_d^* in (2.20) is of the same nature as \mathbf{I}^* in (2.12) because both reduce to computing a complex integral whose integrand is a rational function. In particular, as \mathbf{I}^* , the problem \mathbf{I}_d^* can be solved by Cauchy residue techniques (see for example [14]).

However, there is an important difference between \mathbf{I}^* and \mathbf{I}_d^* . Whereas the data $\{A_{jk}\}$ appears in \mathbf{I}^* as *coefficients* of the dual variables λ_k in $\widehat{F}(\lambda, c)$, it now appears as *exponents* of the dual variables z_k in $\widehat{F}_d(z, c)$. And an immediate consequence of this fact is that the rational function $\widehat{F}_d(\cdot, c)$ has many more poles than $\widehat{F}(\cdot, c)$ (by considering one variable at a time), and in

particular, many of them are *complex*, whereas $\widehat{F}(\cdot, c)$ has only *real* poles. As a result, the integration of $\widehat{F}_d(z, c)$ is more complicated than that of $\widehat{F}(\lambda, c)$, which is reflected in the *discrete* (or *periodic*) Brion and Vergne formula described below. However, we will see that the poles of $\widehat{F}_d(z, c)$ are simply related to those of $\widehat{F}(\lambda, c)$.

2.3.4 The “discrete” Brion and Vergne formula

Brion and Vergne [7] consider the *generating function* $H : \mathbf{C}^m \rightarrow \mathbf{C}$ defined by

$$\lambda \mapsto H(\lambda, c) := \sum_{y \in \mathbf{Z}^m} \widehat{f}_d(y, c) \varepsilon^{-\langle \lambda, y \rangle},$$

which, after the change of variable $z_i = \varepsilon^{\lambda_i}$ for all $i = 1, \dots, m$, reduces to $\widehat{F}_d(z, c)$ in (2.20).

They obtain the nice formula (2.21) below. Namely, and with the same notation used in Section 2.2.4, let $c \in \mathbf{R}^n$ be regular with $-c$ in the interior of $(\mathbf{R}_+^n \cap V)^*$, and let γ be a chamber. Then for all $b \in \Lambda \cap \bar{\gamma}$ (recall $\Lambda = A(\mathbf{Z}^n)$),

$$\widehat{f}_d(b, c) = \sum_{\sigma \in \mathcal{B}(\Delta, \gamma)} \frac{\varepsilon^{c'x(\sigma)}}{\mu(\sigma)} U_\sigma(b, c) \quad (2.21)$$

for some coefficients $U_\sigma(b, c) \in \mathbf{R}$, a detailed expression for which can be found in [7, Theorem 3.4, p. 821]. In particular, due to the occurrence of *complex poles* in $\widehat{F}(z, c)$, the term $U_\sigma(b, c)$ in (2.21) is the *periodic* analogue of $(\prod_{k \notin \sigma} (c_k - \pi_\sigma A_k))^{-1}$ in (2.14).

Again, as for $\widehat{f}(b, c)$, (2.21) can be re-written as

$$\widehat{f}_d(b, c) = \sum_{x(\sigma): \text{vertex of } \Omega(b)} \frac{\varepsilon^{c'x(\sigma)}}{\mu(\sigma)} U_\sigma(b, c), \quad (2.22)$$

to compare with (2.14). To be more precise, by inspection of Brion and Vergne’s formula in [7, p. 821] in our current context, one may see that

$$U_\sigma(b, c) = \sum_{g \in G(\sigma)} \frac{\varepsilon^{2i\pi b(g)}}{V_\sigma(g, c)}, \quad (2.23)$$

where $G(\sigma) := (\oplus_{j \in \sigma} \mathbf{Z}A_j)^*/\Lambda^*$ (where $*$ denotes the dual lattice); it is a finite abelian group of order $\mu(\sigma)$ and with (finitely many) characters $\varepsilon^{2i\pi b}$ for all $b \in \Lambda$. In particular, writing $A_k = \sum_{j \in \sigma} u_{jk} A_j$ for all $k \notin \sigma$,

$$\varepsilon^{2i\pi A_k}(g) = \varepsilon^{2i\pi \sum_{j \in \sigma} u_{jk} g_j} \quad k \notin \sigma.$$

Moreover,

$$V_\sigma(g, c) = \prod_{k \notin \sigma} \left(1 - \varepsilon^{-2i\pi A_k} (g) \varepsilon^{c_k - \pi^\sigma A_k} \right), \quad (2.24)$$

with A_k, π^σ as in (2.13) (and π^σ rational). Again note the importance of the reduced costs $c_k - \pi^\sigma A_k$ in the expression for $\widehat{F}_d(z, c)$.

2.3.5 The discrete optimization problem \mathbf{P}_d

We are now in a position to see how \mathbf{I}_d^* provides some nice information about the optimal value $f_d(b, c)$ of the discrete optimization problem \mathbf{P}_d .

Theorem 1. *Let $A \in \mathbf{Z}^{m \times n}$, $b \in \mathbf{Z}^m$ and let $c \in \mathbf{Z}^n$ be regular with $-c$ in the interior of $(\mathbf{R}_+^n \cap V)^*$. Let $b \in \overline{\gamma} \cap A(\mathbf{Z}^n)$ and let $q \in \mathbf{N}$ be the least common multiple (l.c.m.) of $\{\mu(\sigma)\}_{\sigma \in \mathcal{B}(\Delta, \gamma)}$.*

If $Ax = b$ has no solution $x \in \mathbf{N}^n$ then $f_d(b, c) = -\infty$, else assume that

$$\max_{x(\sigma): \text{vertex of } \Omega(b)} \left[c'x(\sigma) + \lim_{r \rightarrow \infty} \frac{1}{r} \ln U_\sigma(b, rc) \right],$$

is attained at a unique vertex $x(\sigma)$ of $\Omega(b)$. Then

$$\begin{aligned} f_d(b, c) &= \max_{x(\sigma): \text{vertex of } \Omega(b)} \left[c'x(\sigma) + \lim_{r \rightarrow \infty} \frac{1}{r} \ln U_\sigma(b, rc) \right] \\ &= \max_{x(\sigma): \text{vertex of } \Omega(b)} \left[c'x(\sigma) + \frac{1}{q} (\deg(P_{\sigma b}) - \deg(Q_{\sigma b})) \right] \end{aligned} \quad (2.25)$$

for some real-valued univariate polynomials $P_{\sigma b}$ and $Q_{\sigma b}$.

Moreover, the term $\lim_{r \rightarrow \infty} \ln U_\sigma(b, rc)/r$ or $(\deg(P_{\sigma b}) - \deg(Q_{\sigma b}))/q$ in (2.25) is a sum of certain reduced costs $c_k - \pi^\sigma A_k$ (with $k \notin \sigma$).

For a proof see Section 2.6.1.

Remark 1. Of course, (2.25) is not easy to obtain but it shows that the optimal value $f_d(b, c)$ of \mathbf{P}_d is strongly related to the various complex poles of $\widehat{F}_d(z, c)$. It is also interesting to note the crucial role played by the reduced costs $c_k - \pi^\sigma A_k$ in linear programming. Indeed, from the proof of Theorem 1 the optimal value $f_d(b, c)$ is the value of $c'x$ at some vertex $x(\sigma)$ plus a sum of certain reduced costs (see (2.50) and the form of the coefficients $\alpha_j(\sigma, c)$). Thus, as for the LP problem \mathbf{P} , the optimal value $f_d(b, c)$ of \mathbf{P}_d can be found by inspection of (certain sums of) reduced costs associated with each vertex of $\Omega(b)$.

We next derive an asymptotic result that relates the respective optimal values $f_d(b, c)$ and $f(b, c)$ of \mathbf{P}_d and \mathbf{P} .

Corollary 1. *Let $A \in \mathbf{Z}^{m \times n}$, $b \in \mathbf{Z}^m$ and let $c \in \mathbf{R}^n$ be regular with $-c$ in the interior of $(\mathbf{R}_+^n \cap V)^*$. Let $b \in \gamma \cap \Lambda$ and let $x^* \in \Omega(b)$ be an optimal vertex of \mathbf{P} , that is, $f(b, c) = c'x^* = c'x(\sigma^*)$ for $\sigma^* \in \mathcal{B}(\Delta, \gamma)$, the unique optimal basis of \mathbf{P} . Then for $t \in \mathbf{N}$ sufficiently large,*

$$f_d(tb, c) - f(tb, c) = \lim_{r \rightarrow \infty} \left[\frac{1}{r} \ln U_{\sigma^*}(tb, rc) \right]. \quad (2.26)$$

In particular, for $t \in \mathbf{N}$ sufficiently large, the function $t \mapsto f(tb, c) - f_d(tb, c)$ is periodic (constant) with period $\mu(\sigma^)$.*

For a proof see Section 2.6.2. Thus, when $b \in \gamma \cap \Lambda$ is sufficiently large, say $b = tb_0$ with $b_0 \in \Lambda$ and $t \in \mathbf{N}$, the “max” in (2.25) is attained at the unique optimal basis σ^* of the LP (2.5) (see details in Section 2.6.2).

From Remark 1 it also follows that for sufficiently large $t \in \mathbf{N}$, the optimal value $f_d(tb, c)$ is equal to $f(tb, c)$ plus a certain sum of reduced costs $c_k - \pi^{\sigma^*} A_k$ (with $k \notin \sigma^*$) with respect to the optimal basis σ^* .

2.3.6 A dual comparison of \mathbf{P} and \mathbf{P}_d

We now provide an alternative formulation of Brion and Vergne’s discrete formula (2.22), which explicitly relates dual variables of \mathbf{P} and \mathbf{P}_d . Recall that a *feasible basis* of the linear program \mathbf{P} is a basis $\sigma \in \mathcal{B}(\Delta)$ for which $A_\sigma^{-1}b \geq 0$. Thus let $\sigma \in \mathcal{B}(\Delta)$ be a feasible basis of the linear program \mathbf{P} and consider the system of m equations in \mathbf{C}^m :

$$z_1^{A_{1j}} \cdots z_m^{A_{mj}} = \varepsilon^{c_j}, \quad j \in \sigma. \quad (2.27)$$

Recall that A_σ is the nonsingular matrix $[A_{j_1} | \cdots | A_{j_m}]$, with $j_k \in \sigma$ for all $k = 1, \dots, m$. The above system (2.27) has $\rho(\sigma)$ ($= \det(A_\sigma)$) solutions $\{z(k)\}_{k=1}^{\rho(\sigma)}$, written as

$$z(k) = \varepsilon^\lambda \varepsilon^{2i\pi\theta(k)}, \quad k = 1, \dots, \rho(\sigma) \quad (2.28)$$

for $\rho(\sigma)$ vectors $\{\theta(k)\}$ in \mathbf{C}^m .

Indeed, writing $z = \varepsilon^\lambda \varepsilon^{2i\pi\theta}$ (that is, the vector $\{e^{\lambda_j} \varepsilon^{2i\pi\theta_j}\}_{j=1}^m$ in \mathbf{C}^m) and passing to the logarithm in (2.27) yields

$$A'_\sigma \lambda + 2i\pi A'_\sigma \theta = c_\sigma, \quad (2.29)$$

where $c_\sigma \in \mathbf{R}^m$ is the vector $\{c_j\}_{j \in \sigma}$. Thus $\lambda \in \mathbf{R}^m$ is the unique solution of $A'_\sigma \lambda = c_\sigma$ and θ satisfies

$$A'_\sigma \theta \in \mathbf{Z}^m. \quad (2.30)$$

Equivalently, θ belongs to $(\oplus_{j \in \sigma} A_j \mathbf{Z})^*$, the dual lattice of $\oplus_{j \in \sigma} A_j \mathbf{Z}$.

Thus there is a one-to-one correspondence between the $\rho(\sigma)$ solutions $\{\theta(k)\}$ and the finite group $G'(\sigma) = (\oplus_{j \in \sigma} A_j \mathbf{Z})^* / \mathbf{Z}^m$, where $G(\sigma)$ is a subgroup of $G'(\sigma)$. Thus, with $G(\sigma) = \{g_1, \dots, g_s\}$ and $s := \mu(\sigma)$, we can write $(A'_\sigma)^{-1} g_k = \theta_{g_k} = \theta(k)$, so that for every character $\varepsilon^{2i\pi y}$ of $G(\sigma)$, $y \in \Lambda$, we have

$$\varepsilon^{2i\pi y}(g) = \varepsilon^{2i\pi y' \theta_g}, \quad y \in \Lambda, g \in G(\sigma) \quad (2.31)$$

and

$$\varepsilon^{2i\pi A_j}(g) = \varepsilon^{2i\pi A'_j \theta_g} = 1, \quad j \in \sigma. \quad (2.32)$$

So, for every $\sigma \in \mathcal{B}(\Delta)$, denote by $\{z_g\}_{g \in G(\sigma)}$ these $\mu(\sigma)$ solutions of (2.28), that is,

$$z_g = \varepsilon^\lambda \varepsilon^{2i\pi \theta_g} \in \mathbf{C}^m, \quad g \in G(\sigma), \quad (2.33)$$

with $\lambda = (A'_\sigma)^{-1} c_\sigma$, and where $\varepsilon^\lambda \in \mathbf{R}^m$ is the vector $\{\varepsilon^{\lambda_i}\}_{i=1}^m$.

So, in the linear program \mathbf{P} we have a dual vector $\lambda \in \mathbf{R}^m$ associated with each basis σ . In the integer program \mathbf{P} , with each (same) basis σ there are now associated $\mu(\sigma)$ “dual” (complex) vectors $\lambda + 2i\pi \theta_g$, $g \in G(\sigma)$. Hence, with a basis σ in linear programming, the “dual variables” in integer programming are obtained from (a), the corresponding dual variables $\lambda \in \mathbf{R}^m$ in linear programming, and (b), a periodic correction term $2i\pi \theta_g \in \mathbf{C}^m$, $g \in G(\sigma)$.

We next introduce what we call the *vertex residue function*.

Definition 1. Let $b \in \Lambda$ and let $c \in \mathbf{R}^n$ be regular. Let $\sigma \in \mathcal{B}(\Delta)$ be a feasible basis of the linear program \mathbf{P} and for every $r \in \mathbf{N}$, let $\{z_{gr}\}_{g \in G(\sigma)}$ be as in (2.33), with rc in lieu of c , that is,

$$z_{gr} = \varepsilon^{r\lambda} \varepsilon^{2i\pi \theta_g} \in \mathbf{C}^m; \quad g \in G(\sigma), \quad \text{with } \lambda = (A'_\sigma)^{-1} c_\sigma.$$

The vertex residue function associated with a basis σ of the linear program \mathbf{P} is the function $R_\sigma(z_g, \cdot) : \mathbf{N} \rightarrow \mathbf{R}$ defined by

$$r \mapsto R_\sigma(z_g, r) := \frac{1}{\mu(\sigma)} \sum_{g \in G(\sigma)} \frac{z_{gr}^b}{\prod_{j \notin \sigma} (1 - z_{gr}^{-A_j} \varepsilon^{rc_j})}, \quad (2.34)$$

which is well defined because when c is regular, $|z_{gr}|^{A_k} \neq \varepsilon^{rc_k}$ for all $k \notin \sigma$.

The name *vertex residue* is now clear because in the integration (2.20), $R_\sigma(z_g, r)$ is to be interpreted as a generalized Cauchy residue, with respect to the $\mu(\sigma)$ “poles” $\{z_{gr}\}$ of the generating function $\widehat{F}_d(z, rc)$.

Recall from Corollary 1 that when $b \in \gamma \cap \Lambda$ is sufficiently large, say $b = tb_0$ with $b_0 \in \Lambda$ and some large $t \in \mathbf{N}$, the “max” in (2.25) is attained at the unique optimal basis σ^* of the linear program \mathbf{P} .

Proposition 1. *Let c be regular with $-c \in (\mathbf{R}_+^n \cap V)^*$ and let $b \in \gamma \cap A$ be sufficiently large so that the max in (2.25) is attained at the unique optimal basis σ^* of the linear program \mathbf{P} . Let $\{z_g\}_{g \in G(\sigma^*)}$ be as in (2.33) with $\sigma = \sigma^*$. Then the optimal value of \mathbf{P}_d satisfies*

$$\begin{aligned} f_d(b, c) &= \lim_{r \rightarrow \infty} \frac{1}{r} \ln \left[\frac{1}{\mu(\sigma^*)} \sum_{g \in G(\sigma^*)} \frac{z_{gr}^b}{\prod_{k \notin \sigma^*} (1 - z_{gr}^{-A_k} \varepsilon^{rc_k})} \right] \\ &= \lim_{r \rightarrow \infty} \frac{1}{r} \ln R_{\sigma^*}(z_g, r) \end{aligned} \quad (2.35)$$

and the optimal value of \mathbf{P} satisfies

$$\begin{aligned} f(b, c) &= \lim_{r \rightarrow \infty} \frac{1}{r} \ln \left[\frac{1}{\mu(\sigma^*)} \sum_{g \in G(\sigma^*)} \frac{|z_{gr}|^b}{\prod_{k \notin \sigma^*} (1 - |z_{gr}|^{-A_k} \varepsilon^{rc_k})} \right] \\ &= \lim_{r \rightarrow \infty} \frac{1}{r} \ln R_{\sigma^*}(|z_g|, r). \end{aligned} \quad (2.36)$$

For a proof see Section 2.6.3.

Proposition 1 shows that there is indeed a strong relationship between the integer program \mathbf{P}_d and its continuous analogue, the linear program \mathbf{P} . Both optimal values obey exactly the same formula (2.35), but for the continuous version, the complex vector $z_g \in \mathbf{C}^m$ is replaced by the vector $|z_g| = \varepsilon^{\lambda^*} \in \mathbf{R}^m$ of its component moduli, where $\lambda^* \in \mathbf{R}^m$ is the optimal solution of the LP dual of \mathbf{P} . In summary, when $c \in \mathbf{R}^n$ is regular and $b \in \gamma \cap A$ is sufficiently large, we have the following correspondence.

Linear program \mathbf{P}	Integer program \mathbf{P}_d
unique optimal basis σ^*	unique optimal basis σ^*
1 optimal dual vector $\lambda^* \in \mathbf{R}^m$	$\mu(\sigma^*)$ dual vectors $z_g \in \mathbf{C}^m, g \in G(\sigma^*)$
	$\ln z_g = \lambda^* + 2i\pi\theta_g$
$f(b, c) = \lim_{r \rightarrow \infty} \frac{1}{r} \ln R_{\sigma^*}(z_g , r)$	$f_d(b, c) = \lim_{r \rightarrow \infty} \frac{1}{r} \ln R_{\sigma^*}(z_g, r)$

2.4 A discrete Farkas lemma

In this section we are interested in a discrete analogue of the continuous Farkas lemma. That is, with $A \in \mathbf{Z}^{m \times n}$ and $b \in \mathbf{Z}^m$, consider the issue of the existence of a *nonnegative integral* solution $x \in \mathbf{N}^n$ to the system of linear equations $Ax = b$.

The (continuous) Farkas lemma, which states that given $A \in \mathbf{R}^{m \times n}$ and $b \in \mathbf{R}^m$,

$$\{x \in \mathbf{R}^n \mid Ax = b, x \geq 0\} \neq \emptyset \Leftrightarrow [A' \lambda \geq 0] \Rightarrow b' \lambda \geq 0, \quad (2.37)$$

has no discrete analogue in an *explicit* form. For instance, the *Gomory functions* used in Blair and Jeroslow [6] (see also Schrijver [19, Corollary 23.4b]) are implicitly and iteratively defined, and are not directly defined in terms of the data A, b . On the other hand, for various characterizations of feasibility of the linear diophantine equations $Ax = b$, where $x \in \mathbf{Z}^n$, the interested reader is referred to Schrijver [19, Section 4].

Before proceeding to the general case when $A \in \mathbf{Z}^{m \times n}$, we first consider the case $A \in \mathbf{N}^{m \times n}$, where A (and thus b) has only nonnegative entries.

2.4.1 The case when $A \in \mathbf{N}^{m \times n}$

In this section we assume that $A \in \mathbf{N}^{m \times n}$ and thus necessarily $b \in \mathbf{N}^m$, since otherwise $\{x \in \mathbf{N}^n \mid Ax = b\} = \emptyset$.

Theorem 2. *Let $A \in \mathbf{N}^{m \times n}$ and $b \in \mathbf{N}^m$. Then the following two propositions (i) and (ii) are equivalent:*

- (i) *The linear system $Ax = b$ has a solution $x \in \mathbf{N}^n$.*
- (ii) *The real-valued polynomial $z \mapsto z^b - 1 := z_1^{b_1} \cdots z_m^{b_m} - 1$ can be written*

$$z^b - 1 = \sum_{j=1}^n Q_j(z)(z^{A_j} - 1), \quad (2.38)$$

for some real-valued polynomials $Q_j \in \mathbf{R}[z_1, \dots, z_m]$, $j = 1, \dots, n$, all of which have nonnegative coefficients.

In addition, the degree of the Q_j in (2.38) is bounded by

$$b^* := \sum_{j=1}^m b_j - \min_k \sum_{j=1}^m A_{jk}. \quad (2.39)$$

For a proof see Section 2.6.4. Hence Theorem 2 reduces the issue of existence of a solution $x \in \mathbf{N}^n$ to a particular *ideal membership problem*, that is, $Ax = b$ has a solution $x \in \mathbf{N}^n$ if and only if the polynomial $z^b - 1$ belongs to the *binomial ideal* $I = \langle z^{A_j} - 1 \rangle_{j=1, \dots, n} \subset \mathbf{R}[z_1, \dots, z_m]$ for some weights Q_j with nonnegative coefficients.

Interestingly, consider the ideal $J \subset \mathbf{R}[z_1, \dots, z_m, y_1, \dots, y_n]$ generated by the binomials $z^{A_j} - y_j$, $j = 1, \dots, n$, and let G be a Gröbner basis of J . Using the algebraic approach described in Adams and Loustau [2, Section 2.8], it is known that $Ax = b$ has a solution $x \in \mathbf{N}^n$ if and only if the monomial

z^b can be reduced (with respect to G) to some monomial y^α , in which case, $\alpha \in \mathbf{N}^n$ is a feasible solution. Observe that in this case, we do not know $\alpha \in \mathbf{N}^n$ in advance (we look for it!) to test whether $z^b - y^\alpha \in J$. One has to apply Buchberger's algorithm to (i) find a reduced Gröbner basis G of J , and (ii) reduce z^b with respect to G and check whether the final result is a monomial y^α . Moreover, in the latter approach one uses polynomials in $n + m$ (primal) variables y and (dual) variables z , in contrast with the (only) m dual variables z in Theorem 2.

Remark 2. (a) With b^* as in (2.39) denote by $s(b^*) := \binom{m+b^*}{b^*}$ the dimension of the vector space of polynomials of degree b^* in m variables. In view of Theorem 2, and given $b \in \mathbf{N}^m$, checking the existence of a solution $x \in \mathbf{N}^n$ to $Ax = b$ reduces to checking whether or not there exists a nonnegative solution y to a system of linear equations with:

- $n \times s(b^*)$ variables, the nonnegative coefficients of the Q_j ;
- $s(b^* + \max_k \sum_{j=1}^n A_{jk})$ equations to identify the terms of the same powers on both sides of (2.38).

This in turn reduces to solving an LP problem with $ns(b^*)$ variables and $s(b^* + \max_k \sum_j A_{jk})$ equality constraints. Observe that in view of (2.38), this LP has a matrix of constraints with coefficients made up only of 0's and ± 1 's.

(b) From the proof of Theorem 2 in Section 2.6.4, it easily follows that one may even constrain the weights Q_j in (2.38) to be polynomials in $\mathbf{Z}[z_1, \dots, z_m]$ (instead of $\mathbf{R}[z_1, \dots, z_m]$) with nonnegative coefficients. However, (a) shows that the strength of Theorem 2 is precisely allowing $Q_j \in \mathbf{R}[z_1, \dots, z_m]$ while enabling us to check feasibility by solving a (continuous) linear program. By enforcing $Q_j \in \mathbf{Z}[z_1, \dots, z_m]$ one would end up with an integer linear system whose size was larger than that of the original problem.

2.4.2 The general case

In this section we consider the general case where $A \in \mathbf{Z}^{m \times n}$ so that A may have negative entries, and we assume that the convex polyhedron $\Omega := \{x \in \mathbf{R}_+^n \mid Ax = b\}$ is compact.

The above arguments cannot be repeated because of the occurrence of negative powers. However, let $\alpha \in \mathbf{N}^n$ and $\beta \in \mathbf{N}$ be such that

$$\widehat{A}_{jk} := A_{jk} + \alpha_k \geq 0, \quad k = 1, \dots, n; \quad \widehat{b}_j := b_j + \beta \geq 0, \quad (2.40)$$

for all $j = 1, \dots, m$. Moreover, as Ω is compact, we have that

$$\max_{x \in \mathbf{N}^n} \left\{ \sum_{j=1}^n \alpha_j x_j \mid Ax = b \right\} \leq \max_{x \in \mathbf{R}_+^n} \left\{ \sum_{j=1}^n \alpha_j x_j \mid Ax = b \right\} =: \rho^*(\alpha) < \infty. \quad (2.41)$$

Observe that given $\alpha \in \mathbf{N}^n$, the scalar $\rho^*(\alpha)$ is easily calculated by solving an LP problem. Choose $\mathbf{N} \ni \beta \geq \rho^*(\alpha)$, and let $\widehat{A} \in \mathbf{N}^{m \times n}$ and $\widehat{b} \in \mathbf{N}^m$ be as in (2.40). Then the existence of solutions $x \in \mathbf{N}^n$ to $Ax = b$ is equivalent to the existence of solutions $(x, u) \in \mathbf{N}^n \times \mathbf{N}$ to the system of linear equations

$$\mathbf{Q} \begin{cases} \widehat{A}x + ue_m = \widehat{b} \\ \sum_{j=1}^n \alpha_j x_j + u = \beta. \end{cases} \quad (2.42)$$

Indeed, if $Ax = b$ with $x \in \mathbf{N}^n$ then

$$Ax + e_m \sum_{j=1}^n \alpha_j x_j - e_m \sum_{j=1}^n \alpha_j x_j = b + e_m \beta - e_m \beta,$$

or equivalently,

$$\widehat{A}x + \left(\beta - \sum_{j=1}^n \alpha_j x_j \right) e_m = \widehat{b},$$

and thus, as $\beta \geq \rho^*(\alpha) \geq \sum_{j=1}^n \alpha_j x_j$ (see, for example, (2.41)), we see that (x, u) with $\beta - \sum_{j=1}^n \alpha_j x_j =: u \in \mathbf{N}$ is a solution of (2.42). Conversely, let $(x, u) \in \mathbf{N}^n \times \mathbf{N}$ be a solution of (2.42). Using the definitions of \widehat{A} and \widehat{b} , it then follows immediately that

$$Ax + e_m \sum_{j=1}^n \alpha_j x_j + ue_m = b + \beta e_m; \quad \sum_{j=1}^n \alpha_j x_j + u = \beta,$$

so that $Ax = b$. The system of linear equations (2.42) can be cast in the form

$$B \begin{bmatrix} x \\ u \end{bmatrix} = \begin{bmatrix} \widehat{b} \\ \beta \end{bmatrix} \quad \text{with} \quad B := \begin{bmatrix} \widehat{A} & e_m \\ - & - \\ \alpha' & 1 \end{bmatrix}, \quad (2.43)$$

and as B only has entries in \mathbf{N} , we are back to the case analyzed in Section 2.4.1.

Corollary 2. *Let $A \in \mathbf{Z}^{m \times n}$ and $b \in \mathbf{Z}^m$ and assume that $\Omega := \{x \in \mathbf{R}_+^n \mid Ax = b\}$ is compact. Let $\alpha \in \mathbf{N}^n$ and $\beta \in \mathbf{N}$ be as in (2.40) with $\beta \geq \rho^*(\alpha)$ (see, for example, (2.41)). Then the following two propositions (i) and (ii) are equivalent:*

- (i) *The system of linear equations $Ax = b$ has a solution $x \in \mathbf{N}^n$;*

(ii) *The real-valued polynomial $z \mapsto z^b(z y)^\beta - 1 \in \mathbf{R}[z_1, \dots, z_m, y]$ can be written*

$$z^b(z y)^\beta - 1 = Q_0(z, y)(z y - 1) + \sum_{j=1}^n Q_j(z, y)(z^{A_j}(z y)^{\alpha_j} - 1) \quad (2.44)$$

for some real-valued polynomials $\{Q_j\}_{j=0}^n$ in $\mathbf{R}[z_1, \dots, z_m, y]$, all of which have nonnegative coefficients.

The degree of the Q_j in (2.44) is bounded by

$$(m+1)\beta + \sum_{j=1}^m b_j - \min \left[m+1, \min_{k=1, \dots, n} \left[(m+1)\alpha_k + \sum_{j=1}^m A_{jk} \right] \right].$$

Proof. Let $\widehat{A} \in \mathbf{N}^{m \times n}$, $\widehat{b} \in \mathbf{N}^m$, $\alpha \in \mathbf{N}^n$ and $\beta \in \mathbf{N}$ be as in (2.40) with $\beta \geq \rho^*(\alpha)$. Then apply Theorem 2 to the equivalent form (2.43) of the system Q in (2.42), where B and (\widehat{b}, β) only have entries in \mathbf{N} , and use the definitions of \widehat{A} and \widehat{b} . \square

Indeed Theorem 2 and Corollary 2 have the flavor of a Farkas lemma as it is stated with the transpose A' of A and involving the dual variables z_k associated with the constraints $Ax = b$. In addition, and as expected, it implies the continuous Farkas lemma because if $\{x \in \mathbf{N}^n \mid Ax = b\} \neq \emptyset$, then from (2.44), and with $z := \varepsilon^\lambda$ and $y := (z_1 \cdots z_m)^{-1}$,

$$\varepsilon^{b'\lambda} - 1 = \sum_{j=1}^m Q_j(e^{\lambda_1}, \dots, e^{\lambda_m}, e^{-\sum_i \lambda_i})(\varepsilon^{(A'\lambda)_j} - 1). \quad (2.45)$$

Therefore $A'\lambda \geq 0 \Rightarrow \varepsilon^{(A'\lambda)_j} - 1 \geq 0$ for all $j = 1, \dots, m$, and as the Q_j have nonnegative coefficients, we have $\varepsilon^{b'\lambda} - 1 \geq 0$, which in turn implies $b'\lambda \geq 0$.

Equivalently, evaluating the partial derivatives of both sides of (2.45) with respect to λ_j , at the point $\lambda = 0$, yields $b_j = \sum_{k=1}^n A_{jk} x_k$ for all $j = 1, \dots, m$, with $x_k := Q_k(1, \dots, 1) \geq 0$. Thus $Ax = b$ for some $x \in \mathbf{R}_+^n$.

2.5 Conclusion

We have proposed what we think is a natural duality framework for the integer program \mathbf{P}_d . It essentially relies on the \mathbf{Z} -transform of the associated counting problem \mathbf{I}_d , for which the important Brion and Vergne inverse formula appears to be an important tool for analyzing \mathbf{P}_d . In particular, it shows that the usual *reduced costs* in linear programming, combined with the periodicities phenomena associated with the complex poles of $\widehat{F}_d(z, c)$, also play an essential role for analyzing \mathbf{P}_d . Moreover, for the standard dual

vector $\lambda \in \mathbf{R}^m$ associated with each basis B of the linear program \mathbf{P} , there are $\det(B)$ corresponding dual vectors $z \in \mathbf{C}^m$ for the discrete problem \mathbf{P}_d . Moreover, for b sufficiently large, the optimal value of \mathbf{P}_d is a function of these dual vectors associated with the optimal basis of the linear program \mathbf{P} . A topic of further research is to establish an *explicit* dual optimization problem \mathbf{P}_d^* in these dual variables. We hope that the above results will stimulate further research in this direction.

2.6 Proofs

A proof in French of Theorem 1 can be found in Lasserre [15]. The English proof in [16] is reproduced below.

2.6.1 Proof of Theorem 1

Proof. Use (2.1) and (2.22) to obtain

$$\begin{aligned}
 \varepsilon^{f_d(b,c)} &= \lim_{r \rightarrow \infty} \left[\sum_{x(\sigma): \text{vertex of } \Omega(b)} \frac{\varepsilon^{rc'x(\sigma)}}{\mu(\sigma)} U_\sigma(b, rc) \right]^{1/r} \\
 &= \lim_{r \rightarrow \infty} \left[\sum_{x(\sigma): \text{vertex of } \Omega(b)} \frac{\varepsilon^{rc'x(\sigma)}}{\mu(\sigma)} \sum_{g \in G(\sigma)} \frac{\varepsilon^{2i\pi b(g)}}{V_\sigma(g, rc)} \right]^{1/r} \\
 &= \lim_{r \rightarrow \infty} \left[\sum_{x(\sigma): \text{vertex of } \Omega(b)} H_\sigma(b, rc) \right]^{1/r}. \tag{2.46}
 \end{aligned}$$

Next, from the expression of $V_\sigma(b, c)$ in (2.24), and with rc in lieu of c , we see that $V_\sigma(g, rc)$ is a function of $y := e^r$, which in turn implies that $H_\sigma(b, rc)$ is also a function of ε^r , of the form

$$H_\sigma(b, rc) = (\varepsilon^r)^{c'x(\sigma)} \sum_{g \in G(\sigma)} \frac{\varepsilon^{2i\pi b(g)}}{\sum_j (\delta_j(\sigma, g, A) \times (e^r)^{\alpha_j(\sigma, c)})}, \tag{2.47}$$

for finitely many coefficients $\{\delta_j(\sigma, g, A), \alpha_j(\sigma, c)\}$. Note that the coefficients $\alpha_j(\sigma, c)$ are sums of some reduced costs $c_k - \pi^\sigma A_k$ (with $k \notin \sigma$). In addition, the (complex) coefficients $\{\delta_j(\sigma, g, A)\}$ do not depend on b .

Let $y := \varepsilon^{r/q}$, where q is the l.c.m. of $\{\mu(\sigma)\}_{\sigma \in \mathcal{B}(\Delta, \gamma)}$. As $q(c_k - \pi^\sigma A_k) \in \mathbf{Z}$ for all $k \notin \sigma$,

$$H_\sigma(b, rc) = y^{qc'x(\sigma)} \times \frac{P_{\sigma b}(y)}{Q_{\sigma b}(y)} \tag{2.48}$$

for some polynomials $P_{\sigma b}, Q_{\sigma b} \in \mathbf{R}[y]$. In view of (2.47), the degree of $P_{\sigma b}$ and $Q_{\sigma b}$, which depends on b but *not* on the magnitude of b , is uniformly bounded in b .

Therefore, as $r \rightarrow \infty$,

$$H_\sigma(b, rc) \approx (\varepsilon^{r/q})^{qc'x(\sigma) + \deg(P_{\sigma b}) - \deg(Q_{\sigma b})}, \quad (2.49)$$

so that the limit in (2.46), which is given by $\max_{\sigma} \varepsilon^{c'x(\sigma)} \lim_{r \rightarrow \infty} U_\sigma(b, rc)^{1/r}$ (as we have assumed unicity of the maximizer σ), is also

$$\max_{x(\sigma): \text{ vertex of } \Omega(b)} \varepsilon^{c'x(\sigma) + (\deg(P_{\sigma b}) - \deg(Q_{\sigma b}))/q}.$$

Therefore $f_d(b, c) = -\infty$ if $Ax = b$ has no solution $x \in \mathbf{N}^n$, else

$$f_d(b, c) = \max_{x(\sigma): \text{ vertex of } \Omega(b)} \left[c'x(\sigma) + \frac{1}{q}(\deg(P_{\sigma b}) - \deg(Q_{\sigma b})) \right], \quad (2.50)$$

from which (2.25) follows easily.

2.6.2 Proof of Corollary 1

Proof. Let $t \in \mathbf{N}$ and note that $f(tb, rc) = trf(b, c) = trc'x^* = trc'x(\sigma^*)$. As in the proof of Theorem 1, and with tb in lieu of b , we have

$$\widehat{f}_d(tb, rc)^{\frac{1}{r}} = \varepsilon^{tc'x^*} \left[\frac{U_{\sigma^*}(tb, rc)}{\mu(\sigma^*)} + \sum_{\text{vertex } x(\sigma) \neq x^*} \left[\frac{\varepsilon^{rc'x(\sigma)}}{\varepsilon^{rc'x(\sigma^*)}} \right]^t \frac{U_\sigma(tb, rc)}{\mu(\sigma)} \right]^{\frac{1}{r}}$$

and from (2.47)–(2.48), setting $\delta_\sigma := c'x^* - c'x(\sigma) > 0$ and $y := \varepsilon^{r/q}$,

$$\widehat{f}_d(tb, rc)^{1/r} = \varepsilon^{tc'x^*} \left[\frac{U_{\sigma^*}(tb, rc)}{\mu(\sigma^*)} + \sum_{\text{vertex } x(\sigma) \neq x^*} y^{-tq\delta_\sigma} \frac{P_{\sigma tb}(y)}{Q_{\sigma tb}(y)} \right]^{1/r}.$$

Observe that $c'x(\sigma^*) - c'x(\sigma) > 0$ whenever $\sigma \neq \sigma^*$ because $\Omega(y)$ is simple if $y \in \gamma$, and c is regular. Indeed, as x^* is an optimal vertex of the LP problem \mathbf{P} , the reduced costs $c_k - \pi^{\sigma^*} A_k$ ($k \notin \sigma^*$) with respect to the optimal basis σ^* are all nonpositive, and in fact, strictly negative because c is regular (see Section 2.2.4). Therefore the term

$$\sum_{\text{vertex } x(\sigma) \neq x^*} y^{-tq\delta_\sigma} \frac{P_{\sigma tb}(y)}{Q_{\sigma tb}(y)}$$

is negligible for t sufficiently large, when compared with $U_{\sigma^*}(tb, rc)$. This is because the degrees of P_{σ^*tb} and Q_{σ^*tb} depend on tb but *not* on the magnitude of tb (see (2.47)–(2.48)), and they are uniformly bounded in tb . Hence taking the limit as $r \rightarrow \infty$ yields

$$\varepsilon^{fa(tb,c)} = \lim_{r \rightarrow \infty} \left[\frac{\varepsilon^{rtc'x(\sigma^*)}}{\mu(\sigma^*)} U_{\sigma^*}(tb, rc) \right]^{1/r} = \varepsilon^{tc'x(\sigma^*)} \lim_{r \rightarrow \infty} U_{\sigma^*}(tb, rc)^{1/r},$$

from which (2.26) follows easily.

Finally, the periodicity comes from the term $\varepsilon^{2i\pi tb}(g)$ in (2.23) for $g \in G(\sigma^*)$. The period is then, of the order $G(\sigma^*)$. \square

2.6.3 Proof of Proposition 3.1

Proof. Let $U_{\sigma^*}(b, c)$ be as in (2.23)–(2.24). It follows immediately that $\pi^{\sigma^*} = (\lambda^*)'$ and so

$$\varepsilon^{-\pi^{\sigma^*} A_k} \varepsilon^{-2i\pi A_k}(g) = \varepsilon^{-A'_k \lambda^*} \varepsilon^{-2i\pi A'_k \theta_g} = z_g^{-A_k}, \quad g \in G(\sigma^*).$$

Next, using $c'x(\sigma^*) = b'\lambda^*$,

$$\varepsilon^{c'x(\sigma^*)} \varepsilon^{2i\pi b}(g) = \varepsilon^{b'\lambda^*} \varepsilon^{2i\pi b' \theta_g} = z_g^b, \quad g \in G(\sigma^*).$$

Therefore

$$\begin{aligned} \frac{1}{\mu(\sigma^*)} \varepsilon^{c'x(\sigma)} U_{\sigma^*}(b, c) &= \frac{1}{\mu(\sigma^*)} \sum_{g \in G(\sigma^*)} \frac{z_g^b}{(1 - z_g^{-A_k} \varepsilon^{c_k})} \\ &= R_{\sigma^*}(z_g, 1), \end{aligned}$$

and (2.35) follows from (2.25) because, with rc in lieu of c , z_g becomes $z_{gr} = \varepsilon^{r\lambda^*} \varepsilon^{2i\pi \theta_g}$ (only the modulus changes).

Next, as only the modulus of z_g is involved in (2.36), we have $|z_{gr}| = \varepsilon^{r\lambda^*}$ for all $g \in G(\sigma^*)$, so that

$$\frac{1}{\mu(\sigma^*)} \sum_{g \in G(\sigma^*)} \frac{|z_{gr}|^b}{\prod_{k \notin \sigma^*} (1 - |z_{gr}|^{-A_k} \varepsilon^{rc_k})} = \frac{\varepsilon^{rb'\lambda^*}}{\prod_{k \notin \sigma^*} (1 - \varepsilon^{r(c_k - A'_k \lambda^*)})},$$

and, as $r \rightarrow \infty$,

$$\frac{\varepsilon^{rb'\lambda^*}}{\prod_{k \notin \sigma^*} (1 - \varepsilon^{r(c_k - A'_k \lambda^*)})} \approx \varepsilon^{rb'\lambda^*},$$

because $(c_k - A'_k \lambda^*) < 0$ for all $k \notin \sigma^*$. Therefore

$$\lim_{r \rightarrow \infty} \frac{1}{r} \ln \left[\frac{\varepsilon^{rb' \lambda^*}}{\prod_{k \notin \sigma^*} (1 - \varepsilon^{r(c_k - A'_k \lambda^*)})} \right] = b' \lambda^* = f(b, c),$$

the desired result.

2.6.4 Proof of Theorem 2

Proof. (ii) \Rightarrow (i). Assume that $z^b - 1$ can be written as in (2.38) for some polynomials $\{Q_j\}$ with nonnegative coefficients $\{Q_{j\alpha}\}$, that is, $\{Q_{j(z)}\} = \sum_{\alpha \in \mathbf{N}^m} Q_{j\alpha} z^\alpha = \sum_{\alpha \in \mathbf{N}^m} Q_{j\alpha} z_1^{\alpha_1} \cdots z_m^{\alpha_m}$, for finitely many nonzero (and nonnegative) coefficients $Q_{j\alpha}$. Using the notation of Section 2.3, the function $\widehat{f}_d(b, 0)$, which (as $c = 0$) counts the nonnegative integral solutions $x \in \mathbf{N}^n$ to the equation $Ax = b$, is given by

$$\widehat{f}_d(b, 0) = \frac{1}{(2\pi i)^m} \int_{|z_1|=\gamma_1} \cdots \int_{|z_m|=\gamma_m} \frac{z^{b-e_m}}{\prod_{j=1}^n (1 - z^{-A_k})} dz,$$

where $\gamma \in \mathbf{R}^m$ satisfies $A'\gamma > 0$ (see (2.18) and (2.20)).

Writing z^{b-e_m} as $z^{-e_m}(z^b - 1 + 1)$ we obtain

$$\widehat{f}_d(b, 0) = B_1 + B_2,$$

with

$$B_1 = \frac{1}{(2\pi i)^m} \int_{|z_1|=\gamma_1} \cdots \int_{|z_m|=\gamma_m} \frac{z^{-e_m}}{\prod_{j=1}^n (1 - z^{-A_k})} dz$$

and

$$\begin{aligned} B_2 &:= \frac{1}{(2\pi i)^m} \int_{|z_1|=\gamma_1} \cdots \int_{|z_m|=\gamma_m} \frac{z^{-e_m}(z^b - 1)}{\prod_{j=1}^n (1 - z^{-A_k})} dz \\ &= \sum_{j=1}^n \frac{1}{(2\pi i)^m} \int_{|z_1|=\gamma_1} \cdots \int_{|z_m|=\gamma_m} \frac{z^{A_j - e_m} Q_j(z)}{\prod_{k \neq j} (1 - z^{-A_k})} dz \\ &= \sum_{j=1}^n \sum_{\alpha \in \mathbf{N}^m} \frac{Q_{j\alpha}}{(2\pi i)^m} \int_{|z_1|=\gamma_1} \cdots \int_{|z_m|=\gamma_m} \frac{z^{A_j + \alpha - e_m}}{\prod_{k \neq j} (1 - z^{-A_k})} dz. \end{aligned}$$

From (2.20) (with $b := 0$) we recognize in B_1 the number of solutions $x \in \mathbf{N}^n$ to the linear system $Ax = 0$, so that $B_1 = 1$. Next, again from (2.20) (now with $b := A_j + \alpha$), each term

$$C_{j\alpha} := \frac{Q_{j\alpha}}{(2\pi i)^m} \int_{|z_1|=\gamma_1} \cdots \int_{|z_m|=\gamma_m} \frac{z^{A_j + \alpha - e_m}}{\prod_{k \neq j} (1 - z^{-A_k})} dz$$

is equal to

$$Q_{j\alpha} \times \text{the number of integral solutions } x \in \mathbf{N}^{n-1}$$

of the linear system $\widehat{A}^{(j)}x = A_j + \alpha$, where $\widehat{A}^{(j)}$ is the matrix in $\mathbf{N}^{m \times (n-1)}$ obtained from A by deleting its j -th column A_j . As by hypothesis each $Q_{j\alpha}$ is nonnegative, it follows that

$$B_2 = \sum_{j=1}^n \sum_{\alpha \in \mathbf{N}^m} C_{j\alpha} \geq 0,$$

so that $\widehat{f}_d(b, 0) = B_1 + B_2 \geq 1$. In other words, the system $Ax = b$ has at least one solution $x \in \mathbf{N}^n$.

(i) \Rightarrow (ii). Let $x \in \mathbf{N}^n$ be a solution of $Ax = b$, and write

$$z^b - 1 = z^{A_1 x_1} - 1 + z^{A_1 x_1} (z^{A_2 x_2} - 1) + \dots + z^{\sum_{j=1}^{n-1} A_j x_j} (z^{A_n x_n} - 1)$$

and

$$z^{A_j x_j} - 1 = (z^{A_j} - 1) \left[1 + z^{A_j} + \dots + z^{A_j(x_j-1)} \right], \quad j = 1, \dots, n,$$

to obtain (2.38) with

$$z \mapsto Q_j(z) := z^{\sum_{k=1}^{j-1} A_k x_k} \left[1 + z^{A_j} + \dots + z^{A_j(x_j-1)} \right], \quad j = 1, \dots, n.$$

We immediately see that each Q_j has all its coefficients nonnegative (and even in $\{0, 1\}$).

Finally, the bound on the degree follows immediately from the proof for (i) \Rightarrow (ii). \square

References

1. K. Aardal, R. Weismantel and L. A. Wolsey, Non-standard approaches to integer programming, *Discrete Appl. Math.* **123** (2002), 5–74.
2. W. W. Adams and P. Loustaunau, *An Introduction to Gröbner Bases* (American Mathematical Society, Providence, RI, 1994).
3. F. Bacelli, G. Cohen, G. J. Olsder and J.-P. Quadrat, *Synchronization and Linearity* (John Wiley & Sons, Chichester, 1992).
4. A. I. Barvinok, Computing the volume, counting integral points and exponential sums, *Discrete Comp. Geom.* **10** (1993), 123–141.
5. A. I. Barvinok and J. E. Pommersheim, An algorithmic theory of lattice points in polyhedra, in *New Perspectives in Algebraic Combinatorics*, MSRI Publications **38** (1999), 91–147.
6. C. E. Blair and R. G. Jeroslow, The value function of an integer program, *Math. Programming* **23** (1982), 237–273.

7. M. Brion and M. Vergne, Residue formulae, vector partition functions and lattice points in rational polytopes, *J. Amer. Math. Soc.* **10** (1997), 797–833.
8. J. B. Conway, *Functions of a Complex Variable I*, 2nd ed. (Springer, New York, 1978).
9. D. den Hertog, *Interior Point Approach to Linear, Quadratic and Convex Programming* (Kluwer Academic Publishers, Dordrecht, 1994).
10. O. Güler, Barrier functions in interior point methods, *Math. Oper. Res.* **21** (1996), 860–885.
11. A. Iosevich, Curvature, combinatorics, and the Fourier transform, *Notices Amer. Math. Soc.* **48** (2001), 577–583.
12. A. Khovanskii and A. Pukhlikov, A Riemann-Roch theorem for integrals and sums of quasipolynomials over virtual polytopes, *St. Petersburg Math. J.* **4** (1993), 789–812.
13. J. B. Lasserre and E. S. Zeron, A Laplace transform algorithm for the volume of a convex polytope, *JACM* **48** (2001), 1126–1140.
14. J. B. Lasserre and E. S. Zeron, An alternative algorithm for counting integral points in a convex polytope, *Math. Oper. Res.* **30** (2005), 597–614.
15. J. B. Lasserre, La valeur optimale des programmes entiers, *C. R. Acad. Sci. Paris Ser. I Math.* **335** (2002), 863–866.
16. J. B. Lasserre, Generating functions and duality for integer programs, *Discrete Optim.* **1** (2004), 167–187.
17. G. L. Litvinov, V. P. Maslov and G. B. Shpiz, Linear functionals on idempotent spaces: An algebraic approach, *Dokl. Akad. Nauk.* **58** (1998), 389–391.
18. D. S. Mitrinović, J. Sándor and B. Crstici, *Handbook of Number Theory* (Kluwer Academic Publishers, Dordrecht, 1996).
19. A. Schrijver, *Theory of Linear and Integer Programming* (John Wiley & Sons, Chichester, 1986).
20. V. A. Truong and L. Tunçel, Geometry of homogeneous convex cones, duality mapping, and optimal self-concordant barriers, *Research report COOR #2002-15* (2002), University of Waterloo, Waterloo, Canada.
21. L. A. Wolsey, Integer programming duality: Price functions and sensitivity analysis, *Math. Programming* **20** (1981), 173–195.



<http://www.springer.com/978-0-387-98095-9>

Optimization

Structure and Applications

(Eds.) C.E.M. Pearce; E. Hunt

2009, XVI, 434 p. 21 illus., Hardcover

ISBN: 978-0-387-98095-9