

The Epidemiology of Alcohol and Drug Disorders

Deborah Hasin and Katherine Keyes

Contents

What is Epidemiology?	23
Substance Use in the United States:	
A Historical Overview	24
Alcohol Consumption	24
Drug Use	25
Substance Use in the United States: A Public Health Problem	25
When Does Use Become Pathological?	
Substance Abuse and Dependence	26
Substance Disorders in the	
Diagnostic and Statistical Manual	
of Mental Disorders	27
Substance Disorders: A Categorical	
or Dimensional Trait?	29
Descriptive Epidemiology: The Incidence and Prevalence of Substance Disorders	29
Prevalence and Incidence of Substance Disorders	29
The Course of Substance Disorders	30
Analytic Epidemiology: The Etiology of Substance Disorders	31
Availability—Temporal and Geographical	31
Pricing, Laws, and Advertising	32
Parental and Peer Influences	34
Stress	34
Religiosity	35
Cognition, Personality	35
Subjective Reactions	36
Psychiatric Comorbidity	36
Genetics	37
Conclusion	38
References	38

What is Epidemiology?

The field of epidemiology involves investigation of the distribution and determinants of health conditions in populations or population subgroups. Epidemiological investigations fall under two common domains: descriptive and analytic. Descriptive epidemiologic studies provide estimates of the incidence and prevalence of illnesses or health behaviors. Incidence refers to the proportion of new cases of a particular health outcome during a specific period of time in a specific at-risk population (i.e., among individuals free of the outcome at the beginning of the time period). Prevalence refers to the proportion of a group or population affected with a health condition at a particular point in time. This includes new cases as well as chronic cases that began earlier and continued into the period of observation. Analytic epidemiologic studies focus on identifying causes/risk factors (e.g., genetic variants, contextual circumstances) of illness, often done through retrospective comparison of cases with non-cases or prospective study of disease development among individuals exposed versus unexposed to a particular hypothesized causal factor.

This chapter covers the epidemiology of alcohol and drug abuse and dependence (referred to together as “substance use disorders”). From an epidemiologic standpoint, substance use disorders have common as well as unique characteristics. This chapter identifies common characteristics of the epidemiology of alcohol and drug use disorders, and highlights

D. Hasin (✉)
Departments of Psychiatry and Epidemiology, Columbia University, New York State Psychiatric Institute, New York, NY, USA
e-mail: hasind@nypdrat.cpmc.columbia.edu

some important characteristics unique to specific substances.

Substance Use in the United States: A Historical Overview

Alcohol Consumption

The use of substances to alter mood states has been a part of civilization from pre-historic through modern time periods. Archeological records document the conversion of sugar into fermented beverages for recreational use, as part of religious ceremonies, and as an analgesic or disinfectant as early at 10,000 B.C. [1, 188]. Alcohol remains incorporated into the fabric of many cultures for a variety of uses, including social and recreational use, as a part of religious ceremonies, secular festivities, and as a normative aspect of daily life. Further, moderate consumption is associated with health and longevity, and is considered to be protective against several adverse health outcomes including cardiovascular disease [13].

Long-term historical information on United States alcohol consumption is available through per-capita alcohol consumption statistics derived from sales records. These records show drinking levels in the United States varied greatly over time from the early days of the United States to the twenty first century [169, 172]. Per-capita consumption levels ranged from extraordinarily high levels during the United States colonial period (from an estimated 5.8 gallons per year per capita in 1790 to 7.1 gallons in 1830) to very low levels before and during Prohibition (from an estimated 1.96 gallons in 1916 to 0.97 gallons in 1934). Prohibition refers to the time period during which the United States prohibited the manufacture, sale, and transportation alcoholic beverages were prohibited by the 18th Amendment to the United States Constitution. This period began in 1920, and ended in 1933 with the repeal of the 18th Amendment by the 21st Amendment. From 1935 until 1982, shown in Fig. 1, per-capita alcohol consumption increased steadily to a peak of nearly 2.8 gallons of ethanol per year in 1982 [169]. Since then, consumption has declined, leveling off at about 2.2 gallons of ethanol per year in 1993, and




Fig. 1 Total per-capita ethanol consumption, United States, 1935–2005. Source: Lakins NE, LaVallee RA, Williams GD, Yi H (2007) Surveillance report #82: apparent per capita alcohol consumption: national, state,

and regional trends, 1977–2005. NIAAA, Division of Biometry and Epidemiology, Alcohol Epidemiologic Data System, Rockville, MD, August 2007

remaining at around that level until 2005, with a slight increase from 1999 to 2005. These data are generally consistent with liver cirrhosis mortality statistics, which show similar variations over time [287].

Worldwide, alcohol consumption patterns vary considerably. Consumption is lowest in predominately Muslim countries (e.g., individuals in Afghanistan and Pakistan consume 0.03 and 0.31 l pure alcohol per capita, respectively) and eastern Mediterranean countries, and highest in eastern European countries (e.g., individual in Ukraine and the Russian Federation consume 15.58 and 15.23 l pure alcohol per capita, respectively) and western European countries such as France, Germany, and the United Kingdom [285].

Alcohol consumption is also heterogeneous within countries. For example, about one-third of United States adults do not drink, although per-capita consumption is 9.3 l [216, 224]. Abstainers are rare in Eastern Europe (including Russia and Ukraine), where per-capita consumption, 13.9 l, is the highest in the world [216]. After immigration, immigrants tend to retain the drinking levels of their country of origin rather than changing to the patterns of their new country, for example, Mexican immigrants in the United States [89] and Russian immigrants in Israel [107, 214].

Drug Use

Drugs such as cannabis, opium, and cocaine have been cultivated and used medicinally as well as recreationally for centuries. Opium poppies are believed to have been first grown in the region near modern-day Iraq as early as 3400 B.C. Opium was used primarily as an analgesic and anesthetic, but medical use did not become widespread until the development of the hypodermic needle in the early 1800s [200]. Historical analysis also indicates that marijuana was smoked recreationally and medically in ancient China as early as 2737 B.C. [199]. In South America, societies have grown and consumed coca, the plant grown to create cocaine,

for centuries. The most common mode of administration is to chew the leaves of the coca plant, or to mix the leaves into a tea [252]. In the twentieth century, innovations in pharmacological knowledge led to the development of synthetic drugs such as lysergic acid diethylamide, categorized as a hallucinogen, and methylenedioxymethamphetamine (or “ecstasy”), categorized as an amphetamine.

In Western countries prior to the 1960s, drug use was rare and the few studies that addressed prevalence focused on heroin, with widely varying results [56, 90, 243]. Morphine is believed to have been prescribed often in the nineteenth and early twentieth centuries mainly as a cough suppressant to ease the suffering of individuals with tuberculosis [199], although no data are available to empirically estimate incidence and prevalence. During the Civil War, it is believed that more than 400,000 soldiers became dependent on morphine, as it was liberally prescribed for pain associated battle wounds [199]. More systematic surveys of United States drug use began in the 1960s. A series of national household surveys on drug use conducted by the National Institute on Drug Abuse and later by the Substance Abuse and Mental Health Services Administration showed that illicit drug use, especially marijuana, increased greatly after the late 1960s (Fig. 2). Heroin use also increased in the late 1960s, when the profile of users changed from “bohemians” to inner-city, unemployed males. Yearly surveys of United States youth [140] since 1975 indicate that ~50% of 12th-grade students have used an illicit drug, with a high of 66% in 1982, a low rate of 41% in 1992, and 51% in 2004. Since 1975, over 80% of students felt that marijuana was easily available, ranging from 82.7% in 1992 to 90.4% in 1998.

Substance Use in the United States: A Public Health Problem

While alcohol and drug use is common both in the United States and in many countries worldwide, excess alcohol consumption is estimated to be the 3rd largest cause of United States




Fig. 2 New users of cannabis in the United States, 1965–2002. Source: Substance Abuse and Mental Health Services Administration (2004). Results from the 2003 National Survey on Drug Use and Health: National Findings

preventable mortality [197] and the 5th largest cause of preventable disability worldwide [66]. Excess substance use and substance use disorders are associated with a broad range of adverse outcomes including but not limited to accidents and traffic fatalities [126], domestic violence [25], fetal alcohol syndrome and other pre- and perinatal insults [211, 246], neuropsychological impairment [11], poor medication adherence (e.g., HIV) [229], economic costs and lost productivity [98], psychiatric comorbidity [21, 114], and functional disability [114]. Thus, prevention and intervention of excess substance use is an important public health priority.

When Does Use Become Pathological? Substance Abuse and Dependence

The two major nomenclatures, the *Diagnostic and Statistical Manual of Mental Disorders*, 4th edition and the International Statistical Classification of Diseases and Related Health Problems, 10th Revision, define psychiatric disorders within a common framework for individuals and groups with different training, experience, and interests. Users include medically and behaviorally trained clinicians, neuroscientists, geneticists, investigators conducting

clinical trials, epidemiologists, policy makers, insurance companies and others. Both the *Diagnostic and Statistical Manual of Mental Disorders*, 4th edition and the research version of the International Statistical Classification of Diseases and Related Health Problems, 10th Revision enable diverse groups to arrive at common definitions of disorders by providing specific, generally observable criteria for each disorder. For substance use disorders, the *Diagnostic and Statistical Manual of Mental Disorders*, 4th edition and the International Statistical Classification of Diseases and Related Health Problems, 10th Revision provide diagnostic criteria for two disorders, dependence and abuse (shown in Tables 1 and 2). The *Diagnostic and Statistical Manual of Mental Disorders*, 4th edition and the International Statistical Classification of Diseases and Related Health Problems, 10th Revision also provide symptoms for diagnosing substance-specific intoxication and withdrawal syndromes, and methods for diagnosing substance-induced psychiatric disorders. The *Diagnostic and Statistical Manual of Mental Disorders*, 4th edition was developed in the United States by the American Psychiatric Association and is used in the United States and internationally in research studies. The International Statistical Classification of Diseases and Related Health Problems, 10th Revision was developed by the World

Table 1 Dependence criteria: International statistical classification of diseases and related health problems, 10th revision (ICD-10) and *diagnostic and statistical manual of mental disorders*, 4th edition (DSM-IV)

Substance	ICD-10	DSM-IV
All substances	<p><i>Three or more of the following six symptoms occurring together for at least 1 month, or if less than 1 month, occurring together repeatedly within a 12-month period:</i></p> <ol style="list-style-type: none">1. Tolerance: need for significantly increased amounts of alcohol to achieve intoxication or desired effect or markedly diminished effect with continued use of the same amount of alcohol.2. A physiological withdrawal state of the characteristic withdrawal syndrome for alcohol, or use of alcohol (or closely related substance) to relieve or avoid symptoms.3. Difficulties in controlling drinking in terms of onset, termination, or levels of use: drinking in larger amounts or over a longer period than intended; or a persistent desire or unsuccessful efforts to reduce or control drinking.4. Important alternative pleasures or interests given up or reduced because of drinking; or a great deal of time spent in activities necessary to obtain or use alcohol or to recover from its effects.5. Persisting with drinking despite clear evidence and knowledge of harmful physical or psychological consequences6. A strong desire or sense of compulsion to drink.	<p>A maladaptive pattern of drinking, leading to clinically significant impairment or distress as manifested by <i>three</i> or more of the following seven symptoms occurring in the same 12-month period:</p> <ol style="list-style-type: none">1. Tolerance: need for markedly increased amounts of alcohol to achieve intoxication or desired effect; or markedly diminished effect with continued use of the same amount of alcohol.2. The characteristic withdrawal syndrome for alcohol (or a closely related substance) or drinking to relieve or avoid withdrawal symptoms.3. Persistent desire or one or more unsuccessful efforts to cut down or control drinking.4. Drinking in larger amounts or over a longer period than the person intended.5. Important social, occupational, or recreational activities given up or reduced because of drinking.6. A great deal of time spent in activities necessary to obtain, to use or to recover from the effects of drinking.7. Continued drinking despite knowledge of having a persistent or recurrent physical or psychological problem that is likely to be caused or exacerbated by drinking.

Health Organization and is used internationally, mainly for clinical purposes and governmental reporting.

Substance Disorders in the Diagnostic and Statistical Manual of Mental Disorders

The substance dependence criteria in the *Diagnostic and Statistical Manual of Mental Disorders*, 4th edition, shown in Table 1, are based on the alcohol dependence syndrome [62], which was generalized to drugs in 1981 [286]. Dependence was considered a combination of physiological and psychological processes leading to increasingly impaired control over substance use in the face of negative consequences. Dependence was one “axis” of substance problems, and the consequences of heavy use (social,

legal, medical problems, hazardous use) a different axis of substance problems. This bi-axial concept [61] led to the distinction between abuse criteria (social, role, legal problems or hazardous use, most commonly driving while intoxicated) and dependence (tolerance, withdrawal, numerous indicators of impaired control over use).

The focus on *dependence* is based on its centrality in research and on its psychometric properties. *Diagnostic and Statistical Manual of Mental Disorders*, 4th edition—defined dependence and International Statistical Classification of Diseases and Related Health Problems, 10th Revision—defined dependence have good to excellent reliability across samples and instruments [24, 26, 83, 99, 101, 108, 265], with few exceptions (rare substances; hallucinogens). Dependence validity has also been shown to be good via several study designs. These include: multi-method comparisons [40, 80, 108, 115,

Table 2 Abuse/harmful use criteria: international statistical classification of diseases and related health problems, 10th revision (ICD-10) and *diagnostic and statistical manual of mental disorders*, 4th edition (DSM-IV)

Substance	ICD-10	DSM-IV
All substances	<p>A: Clear evidence that alcohol use contributed to physical or psychological harm, which may lead to disability/adverse consequences.</p> <p>B: The nature of harm should be clearly identifiable (and specified).</p> <p>C: The pattern of use has persisted for at least 1 month or has occurred repeatedly within a 12-month period.</p> <p>D: Symptoms do not meet criteria for any other mental or behavioral disorder related to alcohol in the same time period (except for acute intoxication).</p>	<p>A: Criteria for alcohol dependence have never been met.</p> <p>B: A maladaptive pattern of drinking, leading to clinically significant impairment or distress as manifested by at least <i>one</i> of the following four symptoms occurring within a 12-month period:</p> <ol style="list-style-type: none"> 1. Recurrent use of alcohol resulting in a failure to fulfill major role obligations at work, school, or home (e.g., repeated absences or poor work performance related to alcohol use; alcohol-related absences, suspensions, or expulsions from school; neglect of children or household). 2. Recurrent alcohol use in situations in which it is physically hazardous (e.g., driving an automobile or operating a machine when impaired by alcohol use). 3. Recurrent alcohol-related legal problems (e.g., arrests for alcohol-related disorderly conduct). 4. Continued alcohol use despite having persistent or recurrent social or interpersonal problems caused by or exacerbated by the effects of alcohol (e.g., arguments with spouse about consequences of intoxication).

213, 226, 231]; longitudinal studies [88, 100, 101, 109, 233, 235]; latent variable analysis [16, 97, 201], and construct validation [105, 113]. Animal models of a syndrome of cocaine dependence symptoms (as distinct from use patterns) [50, 222, 266] lend credence to the dependence syndrome not only as a cross-cultural phenomenon, as suggested by a World Health Organization study [40, 101, 213], but a cross-species phenomenon as well.

Substance *abuse* is a different case. Contrary to clinical assumptions, abuse does not necessarily lead to dependence [88, 100, 109, 116, 233, 235]. Further, not all cases of alcohol or drug dependence have abuse symptoms [110, 111]. Dependence is more familial than abuse [103, 109]. *Diagnostic and Statistical Manual of Mental Disorders*, 4th edition—defined alcohol abuse is most often

diagnosed in the general population based on one symptom, driving while intoxicated [104, 106, 156]; preliminary analyses of national data show this is also the case for drug abuse. A *Diagnostic and Statistical Manual of Mental Disorders*, 4th edition—defined diagnosis of abuse may thus depend on the availability of a car, while dependence is a heritable, complex condition.

Various psychometric analyses have been conducted to examine the validity of the Edwards and Gross taxonomy of two distinct, correlated factors for substance abuse and dependence criteria. Confirmatory factor analysis on the alcohol abuse and dependence items has provided mixed evidence; several studies show that a two-factor model best describes abuse and dependence items [84, 97, 201, 202], while several others found evidence of similar model fit for one- and two-factor models, preferring the

one-factor model on the basis of parsimony and high factor correlations [187, 212]. Factor analyses of cannabis abuse and dependence items have generally found support for a one-factor model or similar fit of one- and two-factor models [3, 73, 187, 203, 257], although results from a general population survey support a two-factor model [16]. Taken together, these studies show some support for combining abuse and dependence albeit with some evidence to the contrary. Differences across study may also have occurred due to characteristics of the populations studied (e.g., general population versus community sample, adults versus adolescents). A current unresolved issue for those preferring a single substance use disorder that combines abuse and dependence criteria is a valid threshold for differentiating between cases and non-cases. This issue will need to be resolved if the criteria are to be combined, for example, in the *Diagnostic and Statistical Manual of Mental Disorders*, 5th edition.

Substance Disorders: A Categorical or Dimensional Trait?

Recent psychometric analyses of the substance abuse and dependence criteria have suggested that these disorders are not categorical entities; instead, evidence supports an underlying continuum of alcohol severity across a variety of samples and populations [112, 142, 164, 185, 212, 228]. Such information may be critical when statistical power is limited, as it often is in studies of gene–gene or gene–environment interaction. If *Diagnostic and Statistical Manual of Mental Disorders*, 4th edition—defined alcohol dependence in categorical form is psychometrically sound (i.e., reliable and valid) but dichotomizes an inherently dimensional condition, then converting its elements to a dimensional measure may produce a more informative phenotype for etiologic studies [112]. Future versions of the diagnostic nomenclature will likely incorporate a dimensional form of substance dependence

[122], but further psychometric and etiologic work validating dimensional forms of substance disorders remains necessary.

Descriptive Epidemiology: The Incidence and Prevalence of Substance Disorders

Prevalence and Incidence of Substance Disorders

The most comprehensive epidemiologic United States information on the incidence and prevalence of alcohol disorders comes from the National Epidemiologic Survey on Alcohol and Related Conditions, a longitudinal survey of 43,093 respondents aged 18 years and older conducted in 2001–2002 [81, 82, 85] with a 3-year follow-up of 34,653 respondents [82]. The diagnostic interview was the Alcohol Use Disorder and Associated Disabilities Interview Schedule—*Diagnostic and Statistical Manual of Mental Disorders*, 4th edition Version [82], a structured interview for non-clinicians with high reliability and validity for substance use disorders [26, 83, 108, 227, 265].

In the National Epidemiologic Survey on Alcohol and Related Conditions, the prevalence of current (past 12 months) alcohol abuse and dependence was 4.7 and 3.8%, respectively, for a total prevalence of 8.5% for any current alcohol use disorder [114]. The prevalence of lifetime alcohol abuse and dependence was 17.8 and 12.5%, respectively, for a total prevalence of 30.3% for any lifetime alcohol use disorder [114]. Current and lifetime alcohol disorders are more prevalent in men (current: 12.4%, lifetime: 42.0%) than in women (current: 4.9%, lifetime: 19.5%). Compared with individuals of White race/ethnicity, among whom the current and lifetime prevalence of alcohol disorders was 8.9 and 34.1%, respectively, Blacks, Hispanics, and Asians have a lower prevalence of current and lifetime alcohol disorders (6.9 and 20.6% for current and lifetime alcohol disorders among

Blacks, 7.9 and 21.0% Hispanics, and 4.5 and 11.6% among Asians. Alcohol disorder prevalence is inversely related to age; those in younger age groups are most likely to have an alcohol disorder, with mean ages at onset of alcohol abuse and dependence at 22.5 and 21.9, respectively [114]. The incidence of alcohol dependence was 1.66 per 100 person-years [85], meaning 1.66 cases per year of alcohol dependence for every 100 individuals without alcohol dependence at the beginning of that year. Incidence of alcohol abuse was slightly lower at 1.03 per 100 person-years [85]. In general, predictors of incidence were similar to predictors of prevalence.

Drug disorders were substantially less common than alcohol disorders. The prevalence of current (past 12 months) drug abuse and dependence was 1.4% and 0.6%, respectively, for a total prevalence of 2% for any current drug use disorder [227]. The prevalence of lifetime drug abuse and dependence was 7.7 and 2.6%, respectively, for a total prevalence of 10.3% for any lifetime drug use disorder [227]. Current and lifetime drug disorders are more prevalent in men (current: 2.8%, lifetime: 13.8%) than in women (current: 1.2%, lifetime: 7.1%). Drug disorder prevalence is inversely related to age; those in younger age groups are most likely to have a drug disorder, with mean ages at onset of drug abuse and dependence at 19 years. There is no consistent trend by race for drug disorders [227]. In the National Epidemiologic Survey on Alcohol and Related Conditions, incidence of drug dependence was estimated at 0.32 per 100 person-years of observation [85]; incidence of drug abuse was slightly lower at 0.28 per 100 person-years. In general, predictors of incidence were similar to predictors of prevalence.

The Course of Substance Disorders

Initiation of alcohol consumption and drug use often occurs during adolescence. Onset of alcohol abuse and dependence is most likely among individuals aged 18–29, although 15% of alcohol dependence cases begin before age 18 [127].

Often, substance disorders are not lifelong conditions. Indeed, a high rate of recovery has been documented in general population samples, even among individuals who have never sought treatment. Studies of alcohol disorders in the general population also show that a high proportion of recovered individuals return to moderate drinking as opposed to abstinence [47, 277]. Data from the National Epidemiologic Survey on Alcohol and Related Conditions indicated that approximately 75% of individuals diagnosed with alcohol dependence at some point in the past did not have a current (i.e., past year) diagnosis, but that only about 20% of these individuals were abstinent from alcohol [47]. However, prospective follow-up of this sample has indicated that low-risk drinking represents a risk factor for relapse to an alcohol disorder compared with abstinence [45]. Longer term prospective follow-up of this general population sample will help to clarify the role of alcohol consumption in recovery from disorder.

The transition to adulthood represents a key developmental phase in which alcohol disorders often remit, in a process termed “maturing out” [8, 46]. Major predictors of recovery include key lifestyle components, such as employment, marriage, and childbirth. Whether or not these factors have a causal influence on recovery or reflect common factors underlying the positive lifestyle components and the recovery remains unknown.

Despite substantial progress in the development of treatments for alcohol and drug disorders, only about one-fifth of those individuals with an alcohol disorder [34, 114] and one-sixth of individuals with a drug disorder [35] seek treatment for the condition during their lifetime. Further, the delay from onset of disorder to treatment is typically 8–10 years [276]. Finally, in contrast to sharp increases in treatment utilization for disorders such as depression between 1990 and 2003 [153], a corresponding increase in the proportion of individuals seeking treatment for an alcohol and drug disorders did not occur during this period [114].

The path from first use to dependence to treatment also differs by gender. Women who

use alcohol and drugs often start using later than men, have a faster progression from first use to dependence, and enter treatment sooner than men given equal ages of dependence onset [209, 215], although no such differences have been observed for crack-cocaine users [55, 171]. This phenomenon has been termed “telescoping”.

Evidence is accumulating that these well-documented gender differences in the course of alcohol disorders are converging. Studies of adolescent alcohol use have consistently shown a convergence in rates of alcohol and drug use initiation in younger birth cohorts, especially those born after World War II [139, 141]. Further, several genetically informative samples have researched gender differences in *Diagnostic and Statistical Manual of Mental Disorders*, 4th edition—defined alcohol and drug disorders over time, unanimously finding support for such a convergence [128, 220]. Similarly, large, representative cross-sectional studies in the United States support gender convergence in rates of *Diagnostic and Statistical Manual of Mental Disorders*, 4th edition—defined alcohol abuse and dependence [92, 155]. Finally, evidence indicates that the traditional “telescoping” phenomenon whereby women exhibit later onset of drug use and disorder but earlier treatment and shorter course may be diminishing, as women are more closely approximating men in both onset and course of disorder [129]. Searches into the causes of these shifts are ongoing, but this evidence indicates increased social acceptability of alcohol use by women in younger generations [91].

Analytic Epidemiology: The Etiology of Substance Disorders

Substance use disorders have a complex etiology involving genetic and environmental factors. These occur along a continuum ranging from the macro level consisting of broad social influences, to the micro level, consisting of molecular-level influences. These can be thought of as external to internal levels (Fig. 3). In the remainder of this chapter, we address these levels in turn. We begin with macro/external factors, including societal availability and desirability of the substances, geographic and temporal differences, pricing, laws, and advertising. We next consider externally imposed stress. Intermediate-level factors include religiosity, parental and peer social influences. Moving increasingly toward the micro and internal levels, we consider cognitive and personality variables, subjective responses to substances, and specific risk as well as protective genes. We conclude by discussing gene-by-environment interaction, addressing the idea that since etiologic influences work at various levels, a factor at any level may emerge more clearly if other levels are considered conjointly.

Availability—Temporal and Geographical

Political Events

Political events, both local and global, influence the availability of substances and thus the risk




Fig. 3 Factors affecting substance use and substance use disorders

of substance use and dependence. In 2004, for example, religiously motivated attacks on alcohol retailers in Iraq (*BBC World News*, July 22, 2004) reduced the availability of alcohol locally for that region. After the Taliban government fell in Afghanistan in 2001, heroin production in Afghanistan increased greatly [165], coinciding with increased heroin use among American teenagers [255]. Political instability in South American countries such as Bolivia and Colombia, especially in the 1970s, influenced the production of cocaine and increased the availability of cocaine in the United States [251]. Thus, political events at a great geographic distance may influence local substance use availability and patterns of use.

Outlet Density

Counties, cities or states with higher density of alcohol outlets (places where alcohol is sold) have higher alcohol consumption and higher rates of alcohol-related problems, including hospital admissions, pedestrian injury collisions, and crashes and crash fatalities [33, 237, 256, 261, 262]. Ecologic and multilevel analysis controlling for individual level factors indicates that outlet density is related to higher mean group rates of consumption and drinking norms scores and to driving after drinking [93, 237]. Community-based interventions to limit access to alcohol by reducing the density of outlets have been shown to reduce alcohol-related traffic injury and self-reported consumption [130]. While information regarding outlet (“dealer”) density is unavailable for drugs, the vigorous efforts of parents, schools, and law enforcement agencies to keep drug dealers away from schools are consistent with the same idea.

Pricing, Laws, and Advertising

Pricing

Alcohol taxation is the major determinant of state variation in the price of alcohol, and is thus

a government intervention. An inverse relationship exists between state-level price of alcohol and per-capita consumption or adverse consequences of drinking [30]. Further, higher state-level beer tax is associated with lower prevalence of *Diagnostic and Statistical Manual of Mental Disorders*, 4th edition—defined alcohol dependence [123]. Outside the United States, cutting the tax on spirits has been followed by increased per-capita alcohol consumption [120, 223].

Laws and Law Enforcement: Alcohol

Laws and their enforcement also affect consumption patterns. In the United States, the 18th Amendment to the Constitution outlawed the manufacture, transport, and sale of alcohol from 1920 to 1933. Figure 1 shows that in 1935, per-capita ethanol consumption was very low, but increased steadily afterwards, consistent with cirrhosis mortality rates from the same period [171, 287]. Thus, the 18th Amendment achieved its purpose, but was repealed because it was unacceptable to the public. Similar events occurred in the former Soviet Union, an area of very high per-capita alcohol consumption [285]. In the mid-1980s, the government attempted to restrict consumption. The policies were successful in reducing consumption, but so unpopular that they contributed to the downfall of the government and were eventually reversed [240].

More recently in the United States, enforcement of laws related to drinking and driving has been shown to be an important deterrent to alcohol-related crashes and fatalities. These include driver’s license suspensions [268], and lowering the maximum legal blood alcohol concentration among drivers [69, 259, 269]. In addition, stricter driving-under-the-influence laws and their enforcement are consistently related to decreased hazardous use [182] and alcohol-related traffic fatalities [6, 268].

Minimum-age drinking laws influence the availability and acceptability of consumption among young people. Laws vary considerably by country both in scope and in minimum age [285]. For example, the minimum consumption

age in the United States is 21, while in Cyprus it is 12. Israel did not have a minimum legal drinking age until 2004, but public concern about increased risky drinking among young adults led to the establishment of a national minimum drinking age (18 years) at that time [248]. Some countries have separate age restrictions for consumption and purchase. For example, in Greece the minimum consumption age is 14 while the minimum purchase age is 17. In Italy, there is no age restriction on consumption in private, but a minimum age requirement of 16 to drink in public.

Minimum drinking age laws have a positive effect on community health as well as the health and safety of adolescents. Research in the United States and other developed countries has indicated that minimum drinking age laws reduce traffic crash and fatality rates [68, 241, 268, 271]; positive effects among adolescents include reducing in alcohol consumption and high risk drinking [206]. Additionally, several studies have documented an association between minimum drinking age laws and a reduction in youth suicide [15, 21].

State Distribution Policies

In the United States, states differ in the ways they control availability of alcohol. Some states exert more control through operation of state alcoholic beverage sales, while others exert less control through the licensing of alcohol outlets. This difference impacts sales and consumption patterns [270]. Compared with “wet” counties, “dry” counties, where alcohol is not sold, have lower rates of alcohol-related accidents, driving-under-the-influence arrests, and cirrhosis mortality [283]. International studies corroborate these findings; in Norway, stringent alcohol regulations, such as mandatory closing on Saturdays, led to lower detoxification admissions [223].

Grass-Roots Efforts

Mothers Against Drunk Driving was started in 1980 by a group of women after a teenage

girl was killed by a repeat-offense drunk driver. Mothers Against Drunk Driving, a very active organization, national since the early 1980s, has been highly effective in influencing state legislation pertaining to intoxicated driving, such as increasing the minimum drinking age from 18 to 21, and enforcement of maximum-blood-alcohol-level laws among drivers [95]. In particular, a highly publicized media campaign called “Rate the State” in which states were graded A through D on driving-under-the-influence countermeasures, put pressure on legislators to increase the stringency of these laws, shown as an effective strategy in reducing alcohol-impaired driving [242, 267].

Alcohol Marketing and Advertising

Product development and marketing aim to increase sales and consumption [29]. Alcohol companies allocate substantial resources to researching consumer preferences, developing new products and promoting them [138]. For example, the alcohol beverage industry spent 696 million dollars on magazine advertising alone between 1997 and 2001, largely targeted to adolescents [74]. The alcohol industry does not publish the results of its marketing research, and resources necessary for definitive public health studies of advertising and other marketing effects are limited by comparison.

Public health concerns often focus on marketing that targets adolescents [28, 39]. Existing data from longitudinal studies show associations between late childhood-early adolescent exposure to advertising and subsequent drinking initiation and frequency [37, 64, 249]. Cross-sectional studies also show associations of various marketing and advertising strategies with positive attitudes about drinking and drinking frequency [71, 167]. Further, an imaging study of adolescent response to alcohol advertising indicated greater brain activation in areas linked to reward and desire among adolescents with alcohol use disorders than infrequent drinkers [254], suggesting that advertisements are especially salient to vulnerable adolescents.

Laws and Law Enforcement: Drugs

A literature on government efforts to reduce drug use by reducing availability is inconsistent. Some studies suggest the strategies are ineffective [17, 279, 284], while others find supply reductions efficacious [48, 278]. Reducing the supply of specific drugs can have unintended consequences, including increases in other substances [260]. Data from United States college studies, however, indicate that increased restrictions on alcohol use does not increase marijuana use, as has been hypothesized, but instead serves to decrease both alcohol and marijuana use [281]. Thus, the evidence is inconsistent on the efficacy of government attempts to limit drug use by reducing supply.

Parental and Peer Influences

Parental Modeling of Substance Use

Twin studies indicate that up to half the liability to alcohol dependence is environmental [225]. Parental modeling has been proposed as one such environmental factor affecting subsequent substance use in their children [65]. Adoption studies do not support this, however, since rates of alcoholism in adoptive children of alcoholics are not elevated [132]. One etiologic model with empirical support from twin studies posits that influential factors for substance use and the progression to dependence change over time; environmental and social factors mediate the initiation and use of substances in childhood and adolescence, while genetic factors become more influential in the adult substance use and dependence [151].

Parenting Practices

Poor parental monitoring increases association with substance-abusing peers [117], a risk factor for alcohol misuse (see below, peer influences). Harsh, inconsistent parenting predicts earlier initiation of alcohol use, conduct

problems and poor regulatory competencies [166, 217]. On the other hand, warm yet authoritative parenting styles protect adolescents from alcohol problems [207].

Peers

Peer influence is a strong predictor of adolescent drug and alcohol use and problems [143, 250, 272]. Twin studies show that shared environmental influences such as peers have a significant effect on initiation of alcohol and any drug use [157, 219]. Two models have been proposed to explain peer influence on adolescent substance use, social selection, and socialization [144]. The social selection theory proposes that young adolescents selectively “mate” with friends; those children who display deviant behavior as children will be prone to choose deviant friendships in adolescence [70]. This can lead to initiation of drug use (especially marijuana use) and may be a factor in the transition to “heavier” drugs. It has been further proposed that an underlying trait such as sensation seeking (see below) influences both the selection of peers and substance use [53]. In contrast, the socialization theory proposes that adolescents can be influenced to use substances by peers in their environment [49] via modeling, offers, development of expectancies, and social norms [18, 236]. Substance use by older siblings is also associated with individual substance use [23, 75, 136, 190]. Studies that could examine these various environmental effects while controlling for genetic influences are needed to resolve the social selection/causation debate.

Peers may also be protective. Some United States ethnic/immigrant groups use substances less than the norm [89]. Adolescents from these groups with ethnically homogeneous peers encounter less pressure to use substances [22].

Stress

Drug disorders are often preceded and accompanied by disruptive behavior and conduct

problems [168] that have a shared genetic vulnerability with drug disorders [149]. These behaviors evoke negative reactions from the environment, resulting in stressful life events that are not always independent of the individuals, making a causal direction between stress and disease onset difficult to discern. In animal studies where stress can be experimentally applied, cause and effect are clearer, as is also the case in studies of early stressful experiences in humans that antedate the onset of substance use disorders.

Animal Models

In animal studies, the timing of stress relative to normal development can be experimentally manipulated. In adult animals, substance use increases after physical stressors [76, 210] and social stressors [44, 85, 96, 194].

Early life stressors also contribute to drug-using behaviors in animals. Neonatally isolated rats are more likely to acquire stimulant self-administration behaviors [134, 160, 179] and show higher dopamine levels in response to cocaine than handled rats, suggesting that early stress leads to greater cocaine reward [20, 161]. Early-life rearing stressors predict ethanol seeking in primates [10]. Isolated rearing led to increased drinking of morphine solution under various conditions [5, 184]. Recently developed animal models of $\Delta 9$ -tetrahydrocannabinol self-administration [19] may allow similar studies for cannabis.

Early Stressors and Drug Use in Humans

Childhood stressors, including parental separation, neglect and abuse (physical and sexual) are associated with later substance use, problems and dependence [54, 150, 152]. However, most studies failed to control for parental history of substance abuse, a potential confounder given that substance abuse is associated with poor parenting [174]. One informative study showed that among adolescents with a substance-abusing

parent, strong family cohesion (the opposite of neglect) protected against drug problems [133]. Twin studies allow the study of environmental stressors while controlling for genetic influences and have shown that childhood sexual abuse is an environmental risk factor for substance use disorders [147, 204]. Effects of other childhood adversities (e.g., neglect, physical and emotional abuse) have not been examined in twin studies.

Religiosity

Religiosity has been called “one of the more important environmental factors that affect the risk for substance use and dependence” [148]. An inverse relationship between religiosity and drinking is cross-cultural [4, 7, 208]. Longitudinal studies of adolescents, college and professional students show that religiosity protects against later heavy drinking [9, 183]. Religiosity is strongly correlated within twin pairs due to shared environmental effects [148, 158, 263]. Heritability of drinking differs between religious and non-religious twins, an example of gene-environment interaction [159]. In twins studied longitudinally [148], religiosity predicted later drinking more than drinking predicted later religiosity, suggesting that religiosity is more likely to influence drinking than the reverse. These studies indicate that religiosity is largely environmental and protects against alcohol use disorders. Religiosity also protects against drug disorders [31, 198], although this literature is less extensive.

Cognition, Personality

Substance Expectancies and Motivations

Positive substance expectancies constitute an important risk factor for the development of alcohol dependence [78, 238]. For example, alcohol expectancies are considered the beliefs that drinking alcohol will result in decreased

negative emotions or enhanced positive emotions [77, 245]. These expectancies can be derived from parents and peers, and are believed to be environmentally influenced rather than genetically influenced [244]. Motivations for drinking often fall under four main domains: (1) drinking to obtain social rewards or enhance social interactions; (2) drinking to enhance positive mood; (3) drinking to reduce negative mood, and (4) drinking to avoid social rejection and conform to social norms. While individuals with alcohol disorders often rate all motivations highly, reduction of negative affect and enhancement of positive affect have been prospectively associated with heavy use and alcohol and drug disorders [14, 27, 137].

Personality Traits

No single personality trait predicts alcoholism [239], but traits associated with the development of alcohol use disorders include novelty seeking [32] and sensation seeking [186, 289], traits that are often associated [58, 282]. The heritability of sensation seeking is unclear, with some twin studies suggesting that approximately half of the variance can be attributed to genetic factors [119, 121, 135], and another suggesting a much weaker influence of genetic factors [195]. Additional personality traits related to alcohol use disorders, albeit less consistently, are neuroticism/negative emotionality [288], impulsivity/disinhibition [191], and extraversion/sociability [125]. Similar traits have been examined in relation to drug use disorders. For example, research has shown that impulsivity/inhibition is reliably lower among individuals with drug abuse/dependence [38, 190], whereas negative emotionality tends to be higher [253, 282].

Subjective Reactions

Level of response to alcohol indicates the quantity needed to obtain an effect. Individuals with

a low level of response need to drink more to obtain an effect. This is a genetically influenced characteristic associated with enhanced risk for alcohol use disorders [234]. Level of response varies by ethnicity. Several groups at high risk for alcohol use disorders show low response, including children of alcoholics, Native Americans, and Koreans [63, 198, 273], while high response is found among Jews [234], a group with relatively low levels of alcohol disorders [107, 174]. A low level of response predicts later onset of alcohol dependence in young adult males [232], and may contribute to transition from lighter to heavier drinking in individuals in a heavy-drinking environment [230]. Several chromosomal regions have shown suggestive linkage results to level of response [280] and an association with variations in the ADH1B gene (one of the genes that influences metabolism of alcohol in the liver) has been documented [57], but replication is needed.

Subjective reactions can also be characterized by whether they are positive or negative. A stimulating (reinforcing), rather than sedating, effect of alcohol has been identified in moderate/heavy drinkers [131], as well as untreated alcoholics [258]. In contrast, a flushing reaction to alcohol includes unpleasant physical sensations [124], found among Asians. A strong flushing reaction precludes drinking, while moderate flushing protects against alcohol dependence. Individuals also vary in their subjective responses to marijuana, and positive and/or negative responses are moderately heritable [180].

Psychiatric Comorbidity

Individuals with substance use disorders exhibit higher rates of mood, anxiety, and personality disorders as compared with the general population [35, 86, 87, 114, 154, 221]. For example, national surveys indicate that individuals with an alcohol disorder are approximately 3.0 times more likely to be diagnosed with major depression; the association between drug disorders and major depression is even stronger,

with odds ratios around 7.0 [36, 102]. A strong association has also been documented between substance disorders and antisocial personality disorder. The National Epidemiologic Survey on Alcohol and Related Conditions survey estimates that 39.3 and 72.4% of individuals with antisocial personality disorder meet criteria for lifetime drug disorders and alcohol disorders, respectively [79].

The strong and consistent relationships between substance disorders and other psychiatric disorders have prompted etiologic researchers to evaluate evidence for an underlying vulnerability to psychiatric disorder in general. Adult twin studies indicate at least moderate genetic heritability across disorder [146, 149, 182], and recent genetic studies have indicated specific genes associated with the transmission of several psychiatric disorder in general, rather than particular disorders [52, 274]. “Internalizing” and “externalizing” domains have been proposed as a means of organizing individual disorders into larger, more meaningful groups. Internalizing disorders are often characterized by the anxiety and depression domains, whereas externalizing disorders are often characterized by alcohol, drug, and antisocial personality disorders. Research into the validity and utility of broad versus narrow categorizations of disorder has been a major area of psychiatric research for decades [192], and remains ongoing [162, 163].

Genetics

Family and Twin Studies of Alcohol and Drug Dependence

Alcoholism [41, 205] and drug disorders [193] are familial. Genetic epidemiology studies of heritability use twin samples to compare concordance for a disorder between monozygotic (identical) vs. dizygotic (non-identical) twins. In these studies, significantly higher concordance in identical twins, who share 100% of their genes, compared with non-identical twins, who

share only an average of 50% of their genes, indicates genetic heritability for a disorder. Twin studies of alcohol dependence show substantial heritabilities (50–60%) [118, 218]. Heritability estimates from studies of illicit drugs are more variable, perhaps due to more varied phenotypes (use, heavy use, abuse and dependence); for drug dependence, heritability estimates are similar to alcohol dependence [72, 147, 219]. For all substances, environmental factors appear to influence initiation and continuation of use, while genetic factors move individuals from use to dependence. Also, as noted above, environmental and social factors mediate the initiation and use of substances in childhood and adolescence, while genetic factors become more influential in the adult substance use and dependence [151]. Some twin studies investigating shared heritability of dependence on different substances showed high shared genetic variance between substances [149, 264] while other studies suggest that dependence on different classes of drugs is not genetically interchangeable [264]. Molecular genetics studies may be able to clarify these issues.

Genetics in Epidemiology Studies and Gene × Environment Interaction

The last 5 years have seen considerable progress in the genetics field in general, as well as in identifying genes whose variants show replicated results on relationships to the risk for alcohol and drug dependence. Some of the genes involved include those that affect the process of alcohol metabolism in the liver such as alcohol dehydrogenase 4 (*ADH4*), related to both alcohol [60, 94, 175, 177] and drug dependence [175, 177, 178]. Other well-replicated findings on genes related to the risk for substance dependence involve processes linked to neurotransmission. These include genes influencing gamma-aminobutyric acid, the major inhibitory neurotransmitter in the brain. Genetic variants in *GABRA-2* predict alcohol dependence in United States [42, 43, 59], Russian [170], and German [67, 247] samples, and the outcome of a behavioral treatment for

alcohol dependence in a multi-site study [12]. *GABRA-2* variants were also related to the risk for drug dependence [2]. The functioning of muscarinic cholinergic receptors underlies many brain functions, including attention, learning, memory, and cognition, all potentially related to addictive disorders. Genetic variants influencing this process include *CHRM2*, shown to affect the risk for alcohol and drug dependence [51, 176, 275] and related personality traits [60].

Although twin studies show that genetic and environmental factors are both important, few studies have addressed whether the relationship of specific genetic variants to alcohol and drug dependence is modified by environmental circumstances. This type of research question could be addressed by appropriately designed epidemiologic studies that collect DNA as well as interview information on risk factors. Until recently, a limitation on such studies was the need to extract DNA from blood samples, a difficult task in survey research due to many practical considerations. Fortunately, methods have recently become available to collect DNA through the use of saliva samples, making the inclusion of genetic variables much more feasible in epidemiologic research. An example of this approach includes a study showing that being exposed to childhood maltreatment interacted with a gene influencing stress reactions to predict early onset of drinking among adolescents [145]. Additional studies of this type are under way in Israel [248] and are being planned in the United States.

Studying the interaction between certain genes and specific environmental factors has important implications for the prevention and treatment of alcohol and drug use disorders. First, better knowledge in this area may help early identification of individuals who are unlikely to be able to use drugs or alcohol in moderation for early education, additional support or supervision. Second, the knowledge may help identify individuals exposed to particular stressors that would particularly benefit from intervention. Finally, clearer knowledge of the interaction of environmental with genetic effects may suggest new lines of investigation to determine the biological mechanisms of protective or

risk-enhancing environmental events or conditions, which may eventually aid in developing better treatments.

Conclusion

In summary, a number of factors influencing the risk for substance dependence have been identified. Through trans-disciplinary research, epidemiologists and others can work together in the future to address multi-level factors conjointly.

References

1. Acker CJ, Tracy SW (eds) (2004) *Altering American consciousness: essays on the history of alcohol and drug use in the United States, 1800–2000*. University of Massachusetts, New York
2. Agrawal A, Edenberg HJ, Foroud T, Bierut LJ, Dunne G, Hinrichs AL, Nurnberger JI, Crowe R, Kuperman S, Schuckit MA, Begleiter H, Porjesz B, Dick DM (2006) Association of *GABRA2* with drug dependence in the collaborative study of the genetics of alcoholism sample. *Behav Genet* 36(5):640–650
3. Agrawal A, Lynskey MT (2007) Does gender contribute to heterogeneity in criteria for cannabis abuse and dependence? Results from the national epidemiological survey on alcohol and related conditions. *Drug Alcohol Depend* 88(2–3):300–307
4. Aharonovich E, Hasin D, Rahav G, Meydan J, Neumark Y (2001) Differences in drinking patterns among Ashkenazic and Sephardic Israeli adults. *J Stud Alcohol* 62:301–305
5. Alexander BK, Beyerstein BL, Hadaway PF, Coombs RB (1981) Effect of early and later colony housing on oral ingestion of morphine in rats. *Pharmacol Biochem Behav* 15:571–576
6. Asbridge M, Mann RE, Flam-Zalcman R, Stoduto G (2004) The criminalization of impaired driving in Canada: assessing the deterrent impact of Canada's first per se law. *J Stud Alcohol* 65(4):450–459
7. Azaiza F, Shoham M, Bar-Hamburger R, et al (2008) Patterns of psychoactive substance use among Arab secondary school students in Israel. *Subst Use Misuse* 43(11):1489–1506
8. Bachman JG, O'Malley PM, Schulenberg JE, Johnston LD, Bryant AL, Merline AC (2002) The decline of substance use in young adulthood: changes in social activities, roles, and beliefs. Lawrence Erlbaum, Mahwah, NJ

9. Barnes GM, Farrell MP, Banerjee S (1994) Family influences on alcohol abuse and other problem behaviors among black and white adolescents in a general population sample. *J Res Adolesc* 4: 183–201
10. Barr CS, Schwandt ML, Newman TK, Higley JD (2004) The use of adolescent nonhuman primates to model human alcohol intake: neurobiological genetic, and psychological variables. *Ann N Y Acad Sci* 1021:221–233
11. Bates ME, Bowden SC, Barry D (2002) Neurocognitive impairment associated with alcohol use disorders: implications for treatment. *Exp Clin Psychopharmacol* 10:193–212
12. Bauer LO, Covault J, Harel O, Das S, Gelernter J, Anton R, Kranzler HR (2007) Variation in GABRA2 predicts drinking behavior in project MATCH subjects. *Alcohol Clin Exp Res* 31(11):1780–1787
13. Bertelli AA (2007) Wine, research and cardiovascular disease: instructions for use. *Atherosclerosis* 195(2):242–247
14. Beseler CL, Aharonovich E, Keyes KM, Hasin DS (2008) Adult transition from at-risk drinking to alcohol dependence: the relationship of family history and drinking motives. *Alcohol Clin Exp Res* 32(4):607–616
15. Birckmayer J, Hemenway D (1999) Minimum-age drinking laws and youth suicide, 1970–1990. *Am J Pub Health* 89:1365–1368
16. Blanco C, Harford TC, Nunes E, Grant B, Hasin D (2007) The latent structure of marijuana and cocaine use disorders: results from the National Longitudinal Alcohol Epidemiologic Survey (NLAES). *Drug Alcohol Depend* 91(1): 91–96
17. Blumenthal RN, Lorvick J, Kral AH, et al (1999) Collateral damage in the war on drugs: HIV risk behaviors along injection drug users. *Int J Drug Policy* 10:25–38
18. Borsari B, Carey KB (2001) Peer influences on college drinking: a review of the research. *J Subst Abuse* 13:391–424
19. Braida D, Iosue S, Pegorini S, Sala M (2004) Delta9-tetrahydrocannabinol-induced conditioned place preference and intracerebroventricular self-administration in rats. *Eur J Pharmacol* 506: 63–69
20. Brake WG, Zhang TY, Diorio J, Meaney MJ, Gratton A (2004) Influence of early postnatal rearing conditions on mesocorticolimbic dopamine and behavioral responses to psychostimulants and stressors in adult rats. *Eur J Neurosci* 19: 1863–1874
21. Brent DA (1995) Risk factors for adolescent suicide and suicidal behavior: mental and substance abuse disorders, family environmental factors, and life stress. *Suicide Life Threat Behav* 25(suppl):52–63
22. Brook JS, Brook DW, De La Rosa M, Duque LF, Rodriguez E, Montoya ID, Whiteman M (1998) Pathways to marijuana use among adolescents: cultural/ecological, family, peer, and personality influences. *J Am Acad Child Adolesc Psychiatry* 37:759–766
23. Brook JS, Whiteman M, Brook DW, Gordon AS (1991) Sibling influences on adolescent drug use: older brothers on younger brothers. *J Am Acad Child Adolesc Psychiatry* 30(6):958–966
24. Bucholz KK, Hesselbrock VM, Shayka JJ, Nurnberger JI Jr, Schuckit MA, Schmidt I, Reich T (1995) Reliability of individual diagnostic criterion items for psychoactive substance dependence and the impact on diagnosis. *J Stud Alcohol* 56: 500–505
25. Caetano R, Nelson S, Cunradi C (2001) Intimate partner violence, dependence symptoms and social consequences of drinking among white, black and Hispanic couples in the United States. *Am J Addict* 10(suppl):60–69
26. Canino G, Bravo M, Ramírez R, Febo VE, Rubio-Stipec M, Fernández RL, Hasin D (1999) The Spanish Alcohol Use Disorder and Associated Disabilities Interview Schedule (AUDADIS): reliability and concordance with clinical diagnoses in a Hispanic population. *J Stud Alcohol* 60:790–799
27. Carpenter KM, Hasin D (1998) A prospective evaluation of the relationship between reasons for drinking and DSM-IV alcohol-use disorders. *Addict Behav* 23(1):41–46
28. Casswell S (2004) Alcohol brands in young peoples' everyday lives: new developments in marketing. *Alcohol Alcohol* 39:471–476
29. Centers for Disease Control (CDC): point-of-purchase alcohol marketing and promotion by store type—United States, 2000–2001. *MMWR* 52:310–313, 2003
30. Chaloupka FJ, Saffer H, Grossman M (2002) The effects of price on alcohol consumption and alcohol-related problems. *Alcohol Res Health* 26(1):22–34
31. Chen CY, Dormitzer CM, Bejarano J, Anthony JC (2004) Religiosity and the earliest stages of adolescent drug involvement in seven countries of Latin America. *Am J Epidemiol* 159:1180–1188
32. Cloninger CR, Sigvardsson S, Przybeck TR, Svrakic DM (1995) Personality antecedents of alcoholism in a national area probability sample. *Eur Arch Psychiatry Clin Neurosci* 245:239–244
33. Cohen DA, Mason K, Scribner R (2002) The population consumption model, alcohol control practices, and alcohol-related traffic fatalities. *Prev Med* 34(2):187–197
34. Cohen E, Feinn R, Arias A, Kranzler HR (2007) Alcohol treatment utilization: findings from the National Epidemiologic Survey on Alcohol and Related Conditions. *Drug Alcohol Depend* 86(2–3):214–221

35. Compton WM, Thomas YF, Stinson FS, Grant BF (2007) Prevalence, correlates, disability, and comorbidity of DSM-IV drug abuse and dependence in the United States: results from the national epidemiologic survey on alcohol and related conditions. *Arch Gen Psychiatry* 64(5):566–576
36. Compton WM, Thomas YF, Stinson FS, Grant BF (2007) Prevalence, correlates, disability, and comorbidity of DSM-IV drug abuse and dependence in the United States: results from the national epidemiologic survey on alcohol and related conditions. *Arch Gen Psychiatry* 64(5):566–576
37. Connolly GM, Casswell S, Zhang JF, Silva PA (1994) Alcohol in the mass media and drinking by adolescents: a longitudinal study. *Addiction* 89:1255–1263
38. Conway KP, Swendsen JD, Rounsaville BJ, et al (2002) Personality, drug of choice, and comorbid psychopathology among substance abusers. *Drug Alcohol Depend* 65:225–234
39. Cooke E, Hastings G, Wheeler C, Eadie D, Moskalewicz J, Dabrowska K (2004) Marketing of alcohol to young people: a comparison of the UK and Poland. *Eur Addict Res* 10:1–7
40. Cottler LB, Grant BF, Blaine J, Mavreas V, Pull C, Hasin D, Compton WM, Rubio-Stipec M, Mager D (1997) Concordance of DSM-IV alcohol and drug use disorder criteria and diagnoses as measured by AUDADIS-ADR, CIDI and SCAN. *Drug Alcohol Depend* 47:195–205
41. Cotton NS (1979) The familial incidence of alcoholism: a review. *J Stud Alcohol* 40:89–116
42. Covault J, Gelernter J, Hesselbrock V, Nellissery M, Kranzler HR (2004) Allelic and haplotypic association of GABRA2 with alcohol dependence. *Am J Med Genet* 129B:104–109
43. Covault J, Gelernter J, Jensen K, Anton R, Kranzler HR (2008) Markers in the 5'-region of GABRG1 associate to alcohol dependence and are in linkage disequilibrium with markers in the adjacent GABRA2 gene. *Neuropsychopharmacology* 33(4):837–848
44. Covington HE, Miczek KA (2001) Repeated social-defeat stress, cocaine or morphine. Effects on behavioral sensitization and intravenous cocaine self-administration "binges". *Psychopharmacology* 158:388–398
45. Dawson DA, Goldstein RB, Grant RF (2007) Rates and correlates of relapse among individuals in remission from DSM-IV alcohol dependence: a 3-year follow-up. *Alcohol Clin Exp Res* 31:2036–2045
46. Dawson DA, Grant BF, Stinson FS, Chou PS (2006) Estimating the effect of help-seeking on achieving recovery from alcohol dependence. *Addiction* 101:824–834
47. Dawson DA, Grant BF, Stinson FS, et al (2004) Recovery from DSM-IV alcohol dependence: United States, 2001–2002. *Addiction* 100(3):281–292
48. Day C, Topp L, Rouen D, Darke S, Hall W, Dolan K (2003) Decreased heroin availability in Sydney Australia in early 2001. *Addiction* 98:93–95
49. Deater-Deckard K (2001) Annotation: recent research examining the role of peer relationships in the development of psychopathology. *J Child Psychol Psychiatry* 42:565–579
50. Deroche-Gamonet V, Belin D, Piazza VP (2004) Evidence for addiction-like behavior in the rat. *Science* 305:1014–1017
51. Dick DM, Aliev F, Wang JC, Gruzza RA, Schuckit M, Kuperman S, Kramer J, Hinrichs A, Bertelsen S, Budde JP, Hesselbrock V, Porjesz B, Edenberg HJ, Bierut LJ, Goate A (2008) Using dimensional models of externalizing psychopathology to aid in gene identification. *Arch Gen Psychiatry* 65(3):310–318
52. Dick DM, Aliev F, Wang JC, Saccone S, Hinrichs A, Bertelsen S, Budde J, Saccone N, Foroud T, Nurnberger J Jr, Xuei X, Conneally PM, Schuckit M, Almasy L, Crowe R, Kuperman S, Kramer J, Tischfield JA, Hesselbrock V, Edenberg HJ, Porjesz B, Rice JP, Bierut L, Goate A (2008) A Systematic single nucleotide polymorphism screen to fine-map alcohol dependence genes on chromosome 7 identifies association with a novel susceptibility gene ACN9. *Biol Psychiatry* 63(11):1047–1053
53. Donohew RL, Hoyle RH, Clayton RR, Skinner WF, Colon SE, Rice RE (1999) Sensation seeking and drug use by adolescents and their friends: models for marijuana and alcohol. *J Stud Alcohol* 60:622–631
54. Dube SR, Felitti VJ, Dong M, Chapman DP, Giles WH, Anda RF (2003) Childhood abuse, neglect, and household dysfunction and the risk of illicit drug use: the adverse childhood experiences study. *Pediatrics* 111:564–572
55. Dudish SA, Hatsukami DK (1996) Gender differences in crack users who are research volunteers. *Drug Alcohol Depend* 42(1):55–63
56. DuPont R, Greene M (1973) The dynamics of a heroin addiction epidemic. *Science* 181:716–722
57. Duranceaux NC, Schuckit MA, Eng MY, Robinson SK, Carr LG, Wall TL (2006) Associations of variations in alcohol dehydrogenase genes with the level of response to alcohol in non-Asians. *Alcohol Clin Exp Res* 30(9):1470–1478
58. Earleywine M, Finn PR, Peterson JB, Pihl RO (1992) Factor structure and correlates of the tridimensional personality questionnaire. *J Stud Alcohol* 53:233–238
59. Edenberg HJ, Dick DM, Xuei X, Tian H, Almasy L, Bauer LO, Crowe RR, Goate A, Hesselbrock V, Jones K, Kwon J, Li TK, Nurnberger JI Jr, O'Connor SJ, Reich T, Rice J, Schuckit MA, Porjesz B, Foroud T, Begleiter H (2004) Variations

- in GABRA2, encoding the alpha 2 subunit of the GABA(A) receptor, are associated with alcohol dependence and with brain oscillations. *Am J Hum Genet* 74(4):705–714
60. Edenberg HJ, Foroud T (2006) The genetics of alcoholism: identifying specific genes through family studies. *Addict Biol* 11(3–4), 386–396
61. Edwards G (1986) The alcohol dependence syndrome: a concept as stimulus to enquiry. *Br J Addict* 81:171–183
62. Edwards G, Gross M (1976) Alcohol dependence: provisional description of a clinical syndrome. *BMJ* 1:1058–1061
63. Ehlers CL, Garcia-Andrade C, Wall TL, Cloutier D, Phillips E (1999) Electroencephalographic responses to alcohol challenge in Native American Mission Indians. *Biol Psychiatry* 45:776–787
64. Ellickson PL, Collins RL, Hambarsoomians K, McCaffrey DF (2005) Does alcohol advertising promote adolescent drinking? Results from a longitudinal assessment. *Addiction* 100:235–246
65. Ellis DA, Zucker RA, Fitzgerald HE (1997) The role of family influences in development and risk. *Alcohol Res Health* 21:218–226
66. Ezzati M, Lopez AD, Rodgers A, Vander Hoorn S, Murray CJ (2002) Comparative risk assessment collaborating group. Selected major risk factors and global and regional burden of disease. *Lancet* 360(9343):1347–1360
67. Fehr C, Sander T, Tadic A, Lenzen KP, Anghelescu I, Klawe C, Dahmen N, Schmidt LG, Szegedi A (2006) Confirmation of association of the GABRA2 gene with alcohol dependence by subtype-specific analysis. *Psychiatr Genet* 16(1):9–17
68. Fell JC, Fisher DA, Voas RB, Blackman K, Tippetts AS (2008) The relationship of underage drinking laws to reductions in drinking drivers in fatal crashes in the United States. *Accid Anal Prev* 40(4):1430–1440
69. Fell JC, Voas RB (2006) The effectiveness of reducing illegal blood alcohol concentration (BAC) limits for driving: evidence for lowering the limit to .05 BAC. *J Safety Res* 37(3):233–243
70. Fergusson DM, Woodward LJ, Horwood LJ (1999) Childhood peer relationship problems and young people's involvement with deviant peers in adolescence. *J Abnorm Child Psychol* 27:357–369
71. Fleming K, Thorson E, Atkin CK (2004) Alcohol advertising exposure and perceptions: links with alcohol expectancies and intentions to drink or drinking in underaged youth and young adults. *J Health Commun* 9:3–29
72. Fu Q, Heath AC, Bucholz KK, Nelson E, Goldberg J, Lyons MJ, True WR, Jacob T, Tsuang MT, Eisen SA (2002) Shared genetic risk of major depression, alcohol dependence, and marijuana dependence: contribution of antisocial personality disorder in men. *Arch Gen Psychiatry* 59: 1125–1132
73. Fulkerson A, Harrison PA, Beebe TJ (1999) DSM-IV substance abuse and dependence: are there really two dimensions of substance use disorders in adolescents? *Addiction* 94:495–506
74. Garfield CF, Chung PJ, Rathouz PJ (2003) Alcohol advertising in magazines and adolescent readership. *JAMA* 289:2424–2429
75. Gfroerer J (1987) Correlation between drug use by teenagers and drug use by older family members. *Am J Drug Alcohol Abuse* 13(1–2):95–108
76. Goeders NE, Guerin GF (1994) Non-contingent electric footshock facilitates the acquisition of intravenous cocaine self-administration in rats. *Psychopharmacology* 114:63–70
77. Goldman MS, Rather BC (1993) Substance use disorders: cognitive models and architectures. In: Kendall P, Dobson KS (eds) *Psychopathology and cognition*. Academic, Orlando, FL, pp 245–291
78. Goldman MS, Brown SA, Christiansen BA, Smith GT (1991) Alcoholism and memory: broadening the scope of alcohol-expectancy research. *Psychol Bull* 110:137–146
79. Goldstein RB, Compton WM, Pulay AJ, Ruan WJ, Pickering RP, Stinson FS, Grant BF (2007) Antisocial behavioral syndromes and DSM-IV drug use disorders in the United States: results from the national epidemiologic survey on alcohol and related conditions. *Drug Alcohol Depend* 90(2–3):145–158
80. Grant BF (1996) DSM-IV, DSM-III-R, and ICD-10 alcohol and drug abuse/harmful use and dependence, United States, 1992: a nosological comparison. *Alcohol Clin Exp Res* 20:1481–1488
81. Grant BF, Dawson DA, Stinson FS, et al (2003) The Alcohol Use Disorder and Associated Disabilities Interview Schedule-IV (AUDADIS-IV): reliability of alcohol consumption, tobacco use, family history of depression and psychiatric diagnostic modules in a general population sample. *Drug Alcohol Depend* 71:7–16
82. Grant BF, Goldstein RB, Chou SP, et al (2008) Sociodemographic and psychopathologic predictors of first incidence of DSM-IV substance use, mood and anxiety disorders: results from the Wave 2 national epidemiologic survey on alcohol and related conditions. *Mol Psychiatry*. Epub ahead of print
83. Grant BF, Harford TC, Dawson DA, Chou PS, Pickering RP (1995) The alcohol use disorder and associated disabilities interview schedule (AUDADIS): reliability of alcohol and drug modules in a general population sample. *Drug Alcohol Depend* 39:37–44
84. Grant BF, Harford TC, Muthén BO, Yi HY, Hasin DS, Stinson FS (2007) DSM-IV alcohol dependence and abuse: further evidence of validity in the general population. *Drug Alcohol Depend* 86 (2–3):154–166
85. Grant BF, Moore TC, Shepard J, Kaplan K Source and accuracy statement: Wave 1 national

- epidemiologic survey on alcohol and related conditions (NESARC). National institute on alcohol abuse and alcoholism Web Site. <http://www.niaaa.nih.gov>. Accessed 2 July 2005
86. Grant BF, Stinson FS, Dawson DA, Chou SP, Dufour MC, Compton W, Pickering RP, Kaplan K (2004) Prevalence and co-occurrence of substance use disorders and independent mood and anxiety disorders: results from the national epidemiologic survey on alcohol and related conditions. *Arch Gen Psychiatry* 61:807–816
 87. Grant BF, Stinson FS, Dawson DA, Chou SP, Dufour MC, Compton W, Pickering RP, Kaplan K (2004) Prevalence and co-occurrence of substance use disorders and independence mood and anxiety disorders: results from the national epidemiologic survey on alcohol and related conditions. *Arch Gen Psychiatry* 61(8):807–816
 88. Grant BF, Stinson FS, Harford TC (2001) Age at onset of alcohol use and DSM-IV alcohol abuse and dependence: a 12-year follow-up. *J Subst Abuse* 13:493–504
 89. Grant BF, Stinson FS, Hasin DS, Dawson DA, Chou SP, Anderson K (2004) Immigration and lifetime prevalence of DSM-IV psychiatric disorders among Mexican Americans and non-Hispanic whites in the United States: results from the national epidemiologic survey on alcohol and related conditions. *Arch Gen Psychiatry* 61: 1226–1233
 90. Greene MH, Nightingale SL, DuPont RL (1975) Evolving patterns of drug abuse. *Ann Intern Med* 83(3):402–411
 91. Greenfield TK, Room R (1997) Situational norms for drinking and drunkenness: trends in the US adult population, 1979–1990. *Addiction* 92(1):33–47
 92. Gruca RA, Norberg K, Bucholz KK, Bierut LJ (2008) Correspondence between secular changes in alcohol dependence and age of drinking onset among women in the United States. *Alcohol Clin Exp Res* 32(8):1493–1501
 93. Gruenewald PJ, Johnson FW, Trepo AJ (2002) Outlets, drinking and driving: a multilevel analysis of availability. *J Stud Alcohol* 63(4): 460–468
 94. Guindalini C, Scivoletto S, Ferreira RG, Breen G, Zilberman M, Peluso MA, Zatz M (2005) Association of genetic variants in alcohol dehydrogenase 4 with alcohol dependence in Brazilian patients. *Am J Psychiatry* 162(5):1005–1007
 95. Hamilton W (2000) Mothers against drunk driving – MADD in the USA. *Inj Prev* 6:90–91
 96. Haney M, Maccari S, Le Moal M, Simon H, Piazza PV (1995) Social stress increases the acquisition of cocaine self-administration in male and female rats. *Brain Res* 698:46–52
 97. Harford TC, Muthen B (2001) The dimensionality of alcohol abuse and dependence. a multivariate analysis of DSM-IV symptom items in the national longitudinal survey of youth. *J Stud Alcohol* 62:150–157
 98. Harwood R, Fountain D, Livermore G (1998) The economic costs of alcohol and drug abuse in the United States, 1992. National Institute on Alcohol Abuse and Alcoholism and National Institute on Drug Abuse, Bethesda, MD
 99. Hasin D, Carpenter KM, McCloud S, Smith M, Grant BF (1997) The alcohol use disorder and associated disabilities interview schedule (AUDADIS): reliability of alcohol and drug modules in a clinical example. *Drug Alcohol Depend* 44(2–3):133–141
 100. Hasin D, Grant B, Endicott J (1990) The natural history of alcohol abuse: implications for definitions of alcohol use disorders. *Am J Psychiatry* 147:1537–1541
 101. Hasin D, Grant BF, Cottler L, Blaine J, Towle L, Ustün B, Sartorius N (1997) Nosological comparisons of alcohol and drug diagnoses: a multisite, multi-instrument international study. *Drug Alcohol Depend* 47(3):217–226
 102. Hasin D, Nunes E, Meydan J (2004) Comorbidity of alcohol, drug, and psychiatric disorders: epidemiology. In: Kranzler HR, Tinsley JA (eds) *Dual diagnosis and psychiatric treatment: substance abuse and comorbid disorders*. Marcel Dekker, New York, pp 1–34
 103. Hasin D, Paykin A (1999) Alcohol dependence and abuse diagnoses: concurrent validity in a nationally representative sample. *Alcohol Clin Exp Res* 23:144–150
 104. Hasin D, Paykin A (1999) Dependence symptoms but diagnosis: diagnostic ‘orphans’ in a 1992 national sample. *Drug Alcohol Depend* 53(3): 215–222
 105. Hasin D, Paykin A (1999) DSM-IV alcohol abuse: investigation in a sample of at-risk drinkers in the community. *J Stud Alcohol* 60: 180–187
 106. Hasin D, Paykin A (1999) DSM-IV alcohol abuse: investigation in a sample of at-risk drinkers in the community. *J Stud Alcohol* 60: 181–187
 107. Hasin D, Rahav G, Meydan J, Neumark Y (1998) The drinking of earlier and more recent Russian immigrants to Israel: comparison to other Israelis. *J Subst Abuse* 10:341–353
 108. Hasin D, Samet S, Nunes E, Meydan J, Matseane K, Waxman R (2006) Diagnosis of comorbid psychiatric disorders in substance users assessed with the Psychiatric Research Interview for Substance and Mental Disorders for DSM-IV. *Am J Psychiatry* 163(4):689–696
 109. Hasin D, Van Rossem R, McCloud S, Endicott J (1997) Alcohol dependence and abuse diagnoses: validity in community sample heavy drinkers. *Alcohol Clin Exp Res* 21(2):213–219
 110. Hasin DS, Grant BF (2004) The co-occurrence of DSM-IV alcohol abuse in DSM-IV alcohol dependence: NESARC results on heterogeneity that

- differs by population subgroup. *Arch Gen Psychiatry* 61:891–896
111. Hasin DS, Hatzenbuehler M, Smith S, Grant BF (2005) Co-occurring DSM-IV drug abuse in DSM-IV drug dependence: results from the national epidemiologic survey on alcohol and related conditions. *Drug Alcohol Depend* 80(1): 117–123
 112. Hasin DS, Liu X, Alderson D, Grant BF (2006) DSM-IV alcohol dependence: a categorical or dimensional phenotype? *Psychol Med* 36(12):1695–1705
 113. Hasin DS, Schuckit MA, Martin CS, Grant BF, Bucholz KK, Helzer JE (2003) The validity of DSM-IV alcohol dependence: what do we know and what do we need to know? *Alcohol Clin Exp Res* 27:244–252
 114. Hasin DS, Stinson FS, Ogburn E, Grant BF (2007) Prevalence, correlates, disability, and comorbidity of DSM-IV alcohol abuse and dependence in the United States: results from the national epidemiologic survey on alcohol and related conditions. *Arch Gen Psychiatry* 64(7):830–842
 115. Hasin DS, Trautman KD, Miele GM, Samet S, Smith M, Endicott J (1996) Psychiatric research interview for substance and mental disorders: reliability for substance abusers. *Am J Psychiatry* 153:1195–1201
 116. Hasin DS, Van Rossem R, McCloud S, Endicott J (1997) Differentiating DSM-IV alcohol dependence and abuse by course: community heavy drinkers. *J Subst Abuse* 9:127–135
 117. Hawkins J, Catalano RF, Miller JY (1992) Risk and protective factors for alcohol and other drug problems in adolescence and early adulthood: implications for substance abuse prevention. *Psychol Bull* 112:64–105
 118. Heath AC (1995) Genetic influences on alcoholism risk: a review of adoption and twin studies. *Alcohol Health Res World* 19:166–171
 119. Heath AC, Cloninger CR, Martin NG (1994) Testing a model for the genetic structure of personality: a comparison of the personality systems of Cloninger and Eysenck. *J Pers Soc Psychol* 66:762–775
 120. Heeb J, Gmel G, Zurbrugg C, et al (2003) Changes in alcohol consumption following a reduction in the price of spirits. *Addiction* 98:1433–1446
 121. Heiman N, Stallings MC, Hofer SM, Hewitt JK (2003) Investigating age differences in the genetic and environmental structure of the tridimensional personality questionnaire in later adulthood. *Behav Genet* 33:171–180
 122. Helzer JE, Bucholz KK, Bierut LJ, Regier DA, Schuckit MA, Guth SE (2006) Should DSM-V include dimensional diagnostic criteria for alcohol use disorders? *Alcohol Clin Exp Res* 30: 303–310
 123. Henderson C, Liu X, Diez Roux AV, Link BG, Hasin D (2004) The effects of US state income inequality and alcohol policies on symptoms of depression and alcohol dependence. *Soc Sci Med* 58:565–575
 124. Higuchi S, Matsushita S, Murayama M, Takagi, S, Hayashida M (1995) Alcohol and aldehyde dehydrogenase polymorphisms and the risk for alcoholism. *Am J Psychiatry* 152: 1219–1221
 125. Hill SY, Shen S, Lowers L, Locke J (2000) Factors predicting the onset of adolescent drinking in families at high risk for developing alcoholism. *Biol Psychiatry* 45:265–275
 126. Hingson RW, Heeren T, Edwards EM (2008) Age at drinking onset, alcohol dependence, and their relation to drug use and dependence, driving under the influence of drugs, and motor-vehicle crash involvement because of drugs. *J Stud Alcohol Drugs* 69(2):192–201
 127. Hingson RW, Heeren T, Winter MR (2006) Age at drinking onset and alcohol dependence: age at onset, duration and severity. *Arch Pediatr Adolesc Med* 160(7):739–746
 128. Holdcraft LC, Iacono WG (2002) Cohort effects on gender differences in alcohol dependence. *Addiction* 97:1025–1036
 129. Holdcraft LC, Iacono WG (2004) Cross-generational effects on gender differences in psychoactive drug abuse and dependence. *Drug and Alcohol Dependence* 74:147–158
 130. Holder HD, Gruenewald PJ, Pnacki WR, Treno AJ, Grube JW, Saltz RF, Voas RB, Reynolds R, Davis J, Sanchez L, Gaumont G, Roeper P (2000) Effect of community-based interventions on high-risk drinking and alcohol-related injuries. *JAMA* 284(18):2341–2347
 131. Holdstock L, King AC, DeWit H (2000) Subjective and objective responses to ethanol in moderate/heavy and light social drinkers. *Alcohol Clin Exp Res* 24:789–794
 132. Hopfer CJ, Crowley TJ, Hewitt JK (2003) Review of twin and adoption studies of adolescent substance abuse. *J Am Acad Child Adolesc Psychiatry* 42:710–719
 133. Hopfer CJ, Stallings MC, Hewitt JK, Crowley TJ (2003) Family transmission of marijuana use, abuse, and dependence. *J Am Acad Child Adolesc Psychiatry* 42:834–841
 134. Hu M, Crombag HS, Robinson TE, Becker JB (2004) Biological basis of sex differences in the propensity to self-administer cocaine. *Neuropsychopharmacology* 29:81–85
 135. Hur YM, Bouchard TJ (1997) The genetic correlation between impulsivity and sensation seeking traits. *Behav Genet* 27:455–463
 136. Irons DE, McGue M, Iacono WG, Oetting WS (2007) Mendelian randomization: a novel test of the gateway hypothesis and models of gene-environment interplay. *Dev Psychopathol* 19(4):1181–1195
 137. Jackson KM, Sher KJ (2005) Similarities and differences of longitudinal phenotypes across alternate indices of alcohol involvement: a

- methodologic comparison of trajectory approaches. *Psychol Addict Behav* 19:339–351
138. Jackson MC, Hastings G, Wheeler C, Eadie D, Mackintosh AM (2000) Marketing alcohol to young people: implications for industry regulation and research policy. *Addiction* 95(suppl 4): S597–S608
 139. Johnson RA, Gerstein DR (1998) Initiation of use of alcohol, cigarettes, marijuana, cocaine, and other substances in US birth cohorts since 1919. *American J Public Health* 88:27–33
 140. Johnston LD, O'Malley PM, Bachman JG, et al (2003) Monitoring the future: national survey results on drug use, 1975–2003: volume I, secondary school students (NIH Publication No. 04–5507). National Institute on Drug Abuse, Bethesda, MD
 141. Johnston LD, O'Malley PM, Bachman JG, Schulenberg JE (2004) Monitoring the Future national survey results on drug use, 1975–2003. Volume I: secondary school students. Bethesda, MD
 142. Kahler CW, Strong DR (2006) A Rasch model analysis of DSM-IV alcohol abuse and dependence items in the national epidemiological survey on alcohol and related conditions. *Alcohol Clin Exp Res* 30(7):1165–1175
 143. Kandel D (1973) Adolescent marihuana use: the role of parents and peers. *Science* 181:1067–1070
 144. Kandel DB (1985) On processes of peer influences in adolescent drug use: a developmental perspective. *Adv Alcohol Subst Abuse* 4:139–163
 145. Kaufman J, Yang BZ, Douglas-Palumberi H, Crouse-Artus M, Lipschitz D, Krystal JH, Gelernter J (2007) Genetic and environmental predictors of early alcohol use. *Biol Psychiatry* 61(11):1228–1234
 146. Kendler K, Prescott CA (1999) A population-based twin study of lifetime major depression in men and women. *Arch Gen Psychiatry* 56:39–44
 147. Kendler KS, Bulik CM, Silberg J, Hettema JM, Myers J, Prescott CA (2000) Childhood sexual abuse and adult psychiatric and substance use disorders in women: an epidemiological and cotwin control analysis. *Arch Gen Psychiatry* 57:953–959
 148. Kendler KS, Gardner CO, Prescott CA (1997) Religion, psychopathology, and substance use and abuse; a multimeasure, genetic-epidemiologic study. *Am J Psychiatry* 154:322–329
 149. Kendler KS, Jacobson KC, Prescott CA, Neale MC (2003) Specificity of genetic and environmental risk factors for use and abuse/dependence of cannabis, cocaine, hallucinogens, sedatives, stimulants, and opiates in male twins. *Am J Psychiatry* 160:687–695
 150. Kendler KS, Neale MC, Prescott CA, Kessler RC, Heath AC, Corey LA, Eaves LJ (1996) Childhood parental loss and alcoholism in women: a causal analysis using a twin-family design. *Psychol Med* 26:79–95
 151. Kendler KS, Schmitt E, Aggen SH, Prescott CA (2008) Genetic and environmental influences on alcohol, caffeine, cannabis, and nicotine use from early adolescence to middle adulthood. *Arch Gen Psychiatry* 65(6):674–682
 152. Kessler RC, Davis CG, Kendler KS (1997) Childhood adversity and adult psychiatric disorder in the US National Comorbidity Survey. *Psychol Med* 27:1101–1119
 153. Kessler RC, Demler O, Frank RG, et al (2005) Prevalence and treatment of mental disorders, 1990 to 2003. *N Engl J Med* 352(24):2515–2523
 154. Kessler RC, McGonagle KA, Zhao S, Nelson CB, Hughes M, Eshleman S, Wittchen HU, Kendler KS (1994) Lifetime and 12-month prevalence of DSM-III-R psychiatric disorders in the United States. Results from the National Comorbidity Survey. *Arch Gen Psychiatry* 51(1):8–19
 155. Keyes KM, Grant BF, Hasin DS (2008) Evidence for a closing gender gap in alcohol use, abuse, and dependence in the United States population. *Drug Alcohol Depend* 93(1–2):21–29
 156. Keyes KM, Hasin DS (2008) Socio-economic status and problem alcohol use: the positive relationship between income and the DSM-IV alcohol abuse diagnosis. *Addiction* 103(7):1120–1130
 157. Keyes MA, Iacono WG, McGue M (2007) Early onset problem behavior, young adult psychopathology, and contextual risk. *Twin Res Hum Genet* 10(1):45–53
 158. Kirk KM, Maes HH, Neale MC, Heath AC, Martin NG, Eaves LJ (1999) Frequency of church attendance in Australia and the United States: models of family resemblance. *Twin Res* 2: 99–107
 159. Koopmans JR, Slutske WS, van Baal GC, Boomsma DI (1999) The influence of religion on alcohol use initiation: evidence for genotype × environment interaction. *Behav Genet* 29:445–453
 160. Kosten TA, Miserebino MJD, Kehoe P (2000) Enhanced acquisition of cocaine self-administration in adult rats with neonatal isolation stress experience. *Brain Res* 875:44–50
 161. Kosten TA, Zhang XY, Kehoe P (2003) Chronic neonatal isolation stress enhances cocaine-induced increases in ventral striatal dopamine levels in rat pups. *Brain Res Dev Brain Res* 141: 109–116
 162. Kramer MD, Krueger RF, Hicks BM (2008) The role of internalizing and externalizing liability factors in accounting for gender differences in the prevalence of common psychopathological syndromes. *Psychol Med* 38(1), 51–61
 163. Krueger RF, Markon KE, Patrick CJ, Benning SD, Kramer MD (2007) Linking antisocial behavior, substance use, and personality: an integrative quantitative model of the adult externalizing spectrum. *J Abnorm Psychol* 116(4):645–666
 164. Krueger RF, Nichol PE, Hicks BM, Markon KE, Patrick CJ, Iacono WG, McGue M (2004) Using

- latent trait modeling to conceptualize an alcohol problems continuum. *Psychol Assess* 16(2): 107–119
165. Kulsudjarit K (2004) Drug problem in southeast and southwest Asia. *Ann NY Acad Sci* 1025: 446–457
 166. Kumpfer KL, Bluth B (2004) Parent/child transactional processes predictive of resilience or vulnerability to “substance use disorders.” *Subst Use Misuse* 39:671–698
 167. Kuo M, Wechsler H, Greenberg P, Lee H (2003) The marketing of alcohol to college students: the role of low prices and special promotions. *Am J Prev Med* 25:204–211
 168. Kuperman S, Schlosser SS, Kramer JR, Bucholz K, Hesselbrock V, Reich T, Reich W (2001) Developmental sequence from disruptive behavior diagnosis to adolescent alcohol dependence. *Am J Psychiatry* 158:2022–2026
 169. Lakins NE, Williams GD, Yi H, Hilton ME (2006) Surveillance Report #78: apparent per capita alcohol consumption: national, state, and regional trends, 1977–2004. NIAAA, Division of Epidemiology and Prevention Research, Alcohol Epidemiologic Data System, Rockville, MD
 170. Lappalainen J, Krupitsky E, Remizov M, Pchelina S, Taraskina A, Zvartau E, Somberg LK, Covault J, Kranzler HR, Krystal JH, Gelernter J (2005) Association between alcoholism and gamma-amino butyric acid alpha2 receptor subtype in a Russian population. *Alcohol Clin Exp Res* 29(4):493–498
 171. Lejeuz CW, Bornoalova MA, Reynolds EK, Daughters SB, Curtin JJ (2007) Risk factors in the relationship between gender and crack/cocaine. *Experimental and Clinical Psychopharmacol* 15(2):165–175
 172. Lender ME, Martin JK (1982) *Drinking in America: a history*. Free, New York
 173. Levav I, Kohn R, Golding JM, Weissman MM (1997) Vulnerability of Jews to affective disorders. *Am J Psychiatry* 154:941–947
 174. Locke TF, Newcomb M (2004) Child maltreatment, parent alcohol and drug-related problems, polydrug problems, and parenting practices: a test of gender differences and four theoretical perspectives. *J Fam Psychol* 18:120–134
 175. Luo X, Kranzler HR, Zuo L, Lappalainen J, Yang BZ, Gelernter J (2006) ADH4 gene variation is associated with alcohol dependence and drug dependence in European Americans: results from HWD tests and case-control association studies. *Neuropsychopharmacology* 31(5):1085–1095
 176. Luo X, Kranzler HR, Zuo L, Wang S, Blumberg HP, Gelernter J (2005) CHRM2 gene predisposes to alcohol dependence; drug dependence and affective disorders: results from an extended case-control structured association study. *Hum Mol Genet* 14(16):2421–2434
 177. Luo X, Kranzler HR, Zuo L, Yang BZ, Lappalainen J, Gelernter J (2005) ADH4 gene variation is associated with alcohol and drug dependence: results from family controlled and population-structured association studies. *Pharmacogenetics Genomics* 15(11):755–768
 178. Luo X, Zuo L, Kranzler HR, Wang S, Anton RF, Gelernter J (2008) Recessive genetic mode of an ADH4 variant in substance dependence in African-Americans: a model of utility of the HWD test. *Behav Brain Funct* 4(1):42
 179. Lynch WJ, Mangigi LD, Taylor JR (2005) Neonatal isolation stress potentiates cocaine seeking behavior in adult male and female rats. *Neuropsychopharmacology* 30:322–329
 180. Lyons MJ, Toomey R, Meyer JM, Green AI, Eisen SA, Goldberg J, True WR, Tsuang MT (1997) How do genes influence marijuana use? The role of subjective effects. *Addiction* 92:409–417
 181. Lyons MJ, True WR, Eisen SA, Goldberg J, Meyer JM, Faraone SV, Eaves LJ, Tsuang MT (1995) Differential heritability of adult and juvenile antisocial traits. *Arch Gen Psychiatry* 52(11):906–915
 182. Maghsoodloo S, Brown DB, Greathouse PA (1988) Impact of the revision of DUI legislation in Alabama. *Am J Drug Alcohol Abuse* 14(1):97–108
 183. Margulies RZ, Kessler RC, Kandel DB (1977) A longitudinal study of onset of drinking among high-school students. *J Stud Alcohol* 38:897–912
 184. Marks-Kaufman R, Lewis JL (1984) Early housing experience modifies morphine self-administration and physical dependence in adult rats. *Addict Behav* 9:235–243
 185. Martin C, Chung T, Kirisci L, Langenbucher JW (2006) Item response theory analysis of diagnostic criteria for alcohol and cannabis use disorders in adolescents: implications for DSM-V. *J Abnorm Psychol* 115:807–14
 186. Martin CA, Kelly TH, Rayens MK, Brogli BR, Brenzel A, Smith WJ, Omar HA (2002) Sensation seeking, puberty, and nicotine, alcohol, and marijuana use in adolescence. *J Am Acad Child Adolesc Psychiatry* 41:1495–1502
 187. Martin CA, Kelly TH, Rayens MK, et al (2004) Sensation seeking and symptoms of disruptive disorder: association with nicotine, alcohol, and marijuana use in early and mid-adolescence. *Psychol Rep* 94:1075–1082
 188. McGovern PE, Zhang J, Tang J, et al (2004) Fermented beverages of pre- and proto-historic China. *Proc Natl Acad Sci USA* 101(51): 17593–17598
 189. McGue M, Sharma A, Benson P (1996) Parent and sibling influences on adolescent alcohol use and misuse: evidence from a U.S. adoption cohort. *J Stud Alcohol* 57(1):8–18
 190. McGue M, Slutske W, Iacono WG (1999) Personality and substance use disorders: II. Alcoholism versus drug use disorders. *J Consult Clin Psychol* 67:394–404

191. McGue M, Slutske W, Taylor J, Iacono WG (1997) Personality and substance use disorders: effects of gender and alcoholism subtype. *Alcohol Clin Exp Res* 21:513–520
192. McKusick VA (1969) On lumpers and splitters, or the nosology of genetic disease. *Perspect Biol Med* 12:298–312
193. Merikangas KR, Stolar M, Stevens DE, Goulet J, Preisig MA, Fenton B, Zhang H, O'Malley SS, Rounsaville BJ (1998) Familial transmission of substance use disorders. *Arch Gen Psychiatry* 55:973–979
194. Miczek KA, Mutschler NH (1996) Activational effects of social stress on IV cocaine self administration in rats. *Psychopharmacology* 128: 256–264
195. Miles DR, van den Bree MB, Gupman AE, Newlin DB, Glantz MD, Pickens RW (2001) A twin study on sensation seeking, risk taking behavior and marijuana use. *Drug Alcohol Depend* 62: 57–68
196. Miller L, Davies M, Greenwald S (2000) Religiosity and substance use and abuse among adolescents in the national comorbidity survey. *J Am Acad Child Adolesc Psychiatry* 39:1190–1197
197. Mokdad AH, Ford ES, Bowman BA, Nelson DE, Engelgau MM, Vinicor F, Marks JS (2000) Diabetes trends in the US: 1990–1998. *Diabetes Care* 23:1278–1283
198. Monteiro MG, Irwin M, Hauger RL, Schuckit MA (1990) TSH response to TRH and family history of alcoholism. *Biol Psychiatry* 27:905–910
199. Musto DF (1999) *The American disease: origins of narcotic control*. Oxford University, New York, Incorporated
200. Musto DF, Korsmeyer P, Malucci TW (Eds.). (2002) *One hundred years of heroin*. Auburn House, New York
201. Muthen B, Grant B, Hasin D (1993) The dimensionality of alcohol abuse and dependence. factor analysis of DSM-III-R and proposed DSM-IV criteria in the 1988 National Health Interview Survey. *Addiction* 88:1079–1090
202. Muthen BO (1995) Factor analysis of alcohol abuse and dependence symptom items in the 1988 national health interview survey. *Addiction* 90:637–645
203. Nelson CB, Rehm J, Ustun TB, Grant B, Chatterji S (1999) Factor structures for DSM-IV substance disorder criteria endorsed by alcohol, cannabis, cocaine and opiate users: results from the WHO reliability and validity study. *Addiction* 94: 843–855
204. Nelson EC, Heath AC, Madden PA, Cooper ML, Dinwiddie SH, Bucholz KK, Glowinski A, McLaughlin T, Dunne MP, Statham DJ, Martin NG (2002) Association between self-reported childhood sexual abuse and adverse psychosocial outcomes: results from a twin study. *Arch Gen Psychiatry* 59:139–145
205. Nurnberger JI Jr, Wiegand R, Bucholz K, O'Connor S, Meyer ET, Reich T, Rice J, Schuckit M, King, L, Petti T, Bierut L, Hinrichs AL, Kuperman S, Hesselbrock V, Porjesz B (2004) A family study of alcohol dependence: coaggregation of multiple disorders in relatives of alcohol-dependent probands. *Arch Gen Psychiatry* 61:1246–1256
206. O'Malley PM, Wagenaar AC (1991) Effects of minimum drinking age laws on alcohol use, related behaviors and traffic crash involvement among american youth: 1976–1987. *J Studies Alcohol* 52(5):478–491
207. Patock-Peckham JA, Cheong J, Balhorn ME, Nagoshi CT (2001) A social learning perspective: a model of parenting styles, self-regulation, perceived drinking control, and alcohol use and problems. *Alcohol Clin Exp Res* 25: 1284–1292
208. Perkins HW (1987) Parental religion and alcohol use problems as intergenerational predictors of problem drinking among college youth. *J Sci Study Relig* 26:340–357
209. Piazza PV, Deminière JM, Le Moal M, Simon H (1989) Factors that predict individual vulnerability to amphetamine self-administration. *Science* 245:1511–1513
210. Piazza PV, Deminière JM, le Moal M, Simon H (1990) Stress- and pharmacologically-induced behavioral sensitization increases vulnerability to acquisition of amphetamine self-administration. *Brain Res* 514:22–26
211. Potter DA, Moreno A, Luther MF, Eddy CA, Siler-Khodr TM, King TS, Schenken RS (1998) Effect of follicular-phase cocaine administration on menstrual and ovarian cyclicity in rhesus monkeys. *American Journal of Obstetrics and Gynecology* 178:118–125
212. Proudfoot H, Baillie AJ, Teesson M (2006) The structure of alcohol dependence in the community. *Drug Alcohol Depend* 81(1):21–26
213. Pull CB, Saunders JB, Mavreas V, Cottler LB, Grant BF, Hasin DS, Blaine J, Mager D, Ustun BT (1997) Concordance between ICD-10 alcohol and drug use disorder criteria and diagnoses as measured by the AUDADIS-ADR, CIDI and SCAN: results of a cross-national study. *Drug Alcohol Depend* 47:207–216
214. Rahav G, Hasin D, Paykin A (1999) Drinking patterns of recent Russian immigrants and other Israelis: 1995 national survey results. *Am J Public Health* 89(8):1212–1216
215. Randall CL, Roberts JS, Del Boca FK, et al (1999) Telescoping of landmark events associated with drinking: a gender comparison. *J Stud Alcohol* 60(2):252–260
216. Rehm J, Rehn N, Room R, et al (2003) The global distribution of average volume of alcohol consumption and patterns of drinking. *Eur Addict Res* 9:147–156

217. Repetti RL, Taylor SE, Seeman TE (2002) Risky families: family social environments and the mental and physical health of offspring. *Psychol Bull* 128:330–66
218. Rhee SH, Hewitt JK, Young SE, Corley RP, Crowley TJ, Stallings MC (2003) Genetic and environmental influences on substance initiation, use, and problem use in adolescents. *Arch Gen Psychiatry* 60:1256–1264
219. Rhee SH, Hewitt JK, Young SE, Corley RP, Crowley TJ, Stallings MC (2003) Genetic and environmental influences on substance initiation, use, and problem use in adolescents. *Arch Gen Psychiatry* 60:1256–1264
220. Rice JP, Neuman RJ, Saccone NL, Corbett J, Rochberg N, Hesselbrock V, Bucholz KK, McGuffin P, Reich T (2003) Age and birth cohort effects on rates of alcohol dependence. *Alcohol Clin Exp Res* 27:93–99
221. Robins LN, Helzer JE, Weissman MM, Orvaschel H, Gruenberg E, Burke JD Jr, Regier DA (1984) Lifetime prevalence of specific psychiatric disorders in three sites. *Arch Gen Psychiatry* 41(10):949–958
222. Robinson TE (2004) Neuroscience. Addicted rats. *Science* 305:951–953
223. Room R (2004) Effects of alcohol controls: Nordic research traditions. *Drug Alcohol* 23:43–53
224. Room R, Graham K, Rehm J, et al (2003) Drinking and its burden in a global perspective: policy considerations and options. *Eur Addict Res* 9:165–175
225. Rose RJ, Dick DM, Viken RJ, Pulkkinen L, Kaprio J (2001) Drinking or abstaining at age 14?: a genetic epidemiological study. *Alcohol Clin Exp Res* 25:1594–1604
226. Rounsaville BJ, Bryant K, Babor T, Kranzler H, Kadden R (1993) Cross system agreement for substance use disorders: DSM-III-R, DSM-IV, and ICD-10. *Addiction* 88:337–348
227. Ruan WJ, Goldstein RB, Chou SP, Smith SM, Saha TD, Pickering RP, Dawson DA, Huang B, Stinson FS, Grant BF (2008) The alcohol use disorder and associated disabilities interview schedule-IV (AUDADIS-IV): reliability of new psychiatric diagnostic modules and risk factors in a general population sample. *Drug Alcohol Depend* 92(1–3):27–36
228. Saha TD, Chou SP, Grant BF (2006) Toward an alcohol use disorder continuum using item response theory: results from the national epidemiologic survey on alcohol and related conditions. *Psychol Med* 36(7):931–941
229. Samet JH, Horton NJ, Meli S, Freedberg KA, Palepu A (2004) Alcohol consumption and antiretroviral adherence among HIV-infected persons with alcohol problems. *Alcohol Clin Exp Res* 28(4):572–577
230. Schuckit MA (1998) Biological, psychological and environmental predictors of the alcoholism risk: a longitudinal study. *J Stud Alcohol* 59:485–494
231. Schuckit MA, Hesselbrock V, Tipp J, et al (1994) Comparison of DSM-III-R, DSM-IV and ICD-10 substance use disorders diagnoses in 1922 men and women subjects in the COGA study. *Addiction* 89:1629–1638
232. Schuckit MA, Smith TL (1996) An 8-year follow-up of 450 sons of alcoholic and control subjects. *Arch Gen Psychiatry* 53:202–210
233. Schuckit MA, Smith TL (2001) A comparison of correlates of DSM-IV alcohol abuse or dependence among more than 400 sons of alcoholics and controls. *Alcohol Clin Exp Res* 25:1–8
234. Schuckit MA, Smith TL, Kalmijn J (2004) The search for genes contributing to the low level of response to alcohol: patterns of findings across studies. *Alcohol Clin Exp Res* 28:1449–1458
235. Schuckit MA, Smith TL, Landi NA (2000) The 5-year clinical course of high-functioning men with DSM-IV alcohol abuse or dependence. *Am J Psychiatry* 157:2028–2035
236. Schulenberg J, O'Malley PM, Bachman JG, Wadsworth KN, Johnston LD (1996) Getting drunk and growing up: trajectories of frequent binge drinking during the transition to young adulthood. *J Stud Alcohol* 57:289–304
237. Scribner RA, Cohen DA, Fisher W (2000) Evidence of a structural effect for alcohol outlet density: a multilevel analysis. *Alcohol Clin Exp Res* 24(2):188–195
238. Shen S, Locke-Wellman J, Hill SY (2001) Adolescent alcohol expectancies in offspring from families at high risk for developing alcoholism. *J Stud Alcohol* 62:763–772
239. Sher KJ, Grekin ER, Williams NA (2005) The development of alcohol use disorders. *Annu Rev Clin Psychol* 1:493–523
240. Shkolnikov V, Nemtsov A (1997) The anti-alcohol campaign and variations in Russian mortality. In: Bobadilla J, Costello C, Mitchell F (eds) *Premature death in the new independent states*. National Academy, Washington, DC
241. Shults RA, Elder RW, Sleet DA, Nichols JL, Alao MO, Carande-Kulis VG, Zaza S, Sosin DM, Thompson RS Task Force on Community Preventive Services. (2008) Reviews of evidence regarding interventions to reduce alcohol-impaired driving. *Am J Prev Med* 21(4 Suppl): 66–88
242. Shults RA, Sleet DA, Elder RW, Ryan GW, Sehgal M (2002) Association between state level drinking and driving countermeasures and self reported alcohol impaired driving. *Inj Prev* 8(2):106–110
243. Singer M The vitality of mythical numbers. *Public Interest* 4:3–9
244. Slutske WS, Cronk NJ, Sher KJ, Madden PA, Bucholz KK, Heath AC (2002) Genes, environment, and individual differences in alcohol expectancies among female adolescents and young adults. *Psychol Addict Behav* 16: 308–317

245. Smith GT (1994) Psychological expectancy as mediator of vulnerability to alcoholism. *Ann NY Acad Sci* 708:165–171
246. Sokol RJ, Delaney-Black V, Nordstrom B (2003) Fetal Alcohol Spectrum Disorder. *J Am Medical Assoc* 290(22):2996–2999
247. Soyka M, Preuss UW, Hesselbrock V, Zill P, Koller G, Bondy B (2008) GABA-A2 receptor subunit gene (GABRA2) polymorphisms and risk for alcohol dependence. *J Psychiatr Res* 42(3):184–191
248. Spivak B, Frisch A, Maman Z, Aharonovich E, Alderson D, Carr LG, Weizman A, Hasin D (2007) Effect of ADH1B genotype on alcohol consumption in young Israeli Jews. *Alcohol Clin Exp Res* 31(8):1297–1301
249. Stacy AW, Zogg JB, Unger JB, Dent CW (2004) Exposure to televised alcohol ads and subsequent adolescent alcohol use. *Am J Health Behav* 28:498–509
250. Steinberg L, Fletcher A, Darling N (1994) Parental monitoring and peer influences on adolescent substance use. *Pediatrics* 93:1060–1064
251. Streatfeild D (2003) Cocaine: an unauthorized biography. Picador, New York
252. Streatfeild D (2002) Cocaine: a definitive history. Virgin Books, New York
253. Swendsen JD, Conway KP, Rounsaville BJ, Merikangas KR (2002) Are personality traits familial risk factors for substance use disorders? Results of a controlled family study. *Am J Psychiatry* 159:1760–1766
254. Tapert SF, Cheung EH, Brown GG, Frank LR, Paulus MP, Schweinsburg AD, Meloy MJ, Brown SA (2003) Neural response to alcohol stimuli in adolescents with alcohol use disorder. *Arch Gen Psychiatry* 60:727–735
255. Tarabar AF, Nelson LS (2003) The resurgence and abuse of heroin by children in the United States. *Curr Opin Pediatr* 15(2):210–215
256. Tatlow JR, Clapp JD, Hohman MM (2000) The relationship between the geographic density of alcohol outlets and alcohol-related hospital admissions in San Diego County. *J Community Health* 25(1):79–88
257. Teesson M, Lynskey M, Manor B, Baillie A (2002) The structure of cannabis dependence in the community. *Drug Alcohol Depend* 68:255–262
258. Thomas SE, Drobles DJ, Voronin K, Anton RF (2004) Following alcohol consumption, nontreatment-seeking alcoholics report greater stimulation but similar sedation compared with social drinkers. *J Stud Alcohol* 65:330–335
259. Tippetts AS, Voas RB, Fell JC, Nichols JL (2005) A meta-analysis of .08 BAC laws in 19 jurisdictions in the United States. *Accid Anal Prev* 37(1):149–161
260. Topp L, Day C, Degenhardt L (2003) Changes in patterns of drug injection concurrent with a sustained reduction in the availability of heroin in Australia. *Drug Alcohol Depend* 70:275–286
261. Treno AJ, Gruenewald PJ, Johnson FW (2001) Alcohol availability and injury: the role of local outlet densities. *Alcohol Clin Exp Res* 92(1–3):27–36
262. Treno AJ, Johnson FW, Remer LG, Gruenewald PJ (2007) The impact of outlet densities on alcohol-related crashes: a spatial panel approach. *Accid Anal Prev* 39(5):894–901
263. Truett KR, Eaves LJ, Walters EE, Heath AC, Hewitt JK, Meyer JM, Silberg J, Neale MC, Martin NG, Kendler KS (1994) A model system for analysis of family resemblance in extended kinships of twins. *Behav Genet* 24:35–49
264. Tsuang MT, Lyons MJ, Meyer JM (1998) Co-occurrence of abuse of different drugs in men: the role of drug-specific and shared vulnerabilities. *Arch Gen Psychiatry* 55:967–972
265. Ustün B, Compton W, Mager D, Babor T, Baiyewu O, Chatterji S, Cottler L, Göğüş A, Mavreas V, Peters L, Pull C, Saunders J, Smeets R, Stipek MR, Vrásti R, Hasin D, Room R, Van den Brink W, Regier D, Blaine J, Grant BF, Sartorius N (1997) WHO Study on the reliability and validity of the alcohol and drug use disorder instruments: overview of methods and results. *Drug Alcohol Depend* 47:161–169
266. Vanderschuren LJ, Everitt BJ (2004) Drug seeking becomes compulsive after prolonged cocaine self-administration. *Science* 305:1017–1019
267. Voas RB, Tippetts AS, Fell J (2000) The relationship of alcohol safety laws to drinking drivers in fatal crashes. *Accid Anal Prev* 32:483–492
268. Voas RB, Tippetts AS, Fell JC (2003) Assessing the effectiveness of minimum legal drinking age and zero tolerance laws in the United States. *Accid Anal Prev* 35(4):579–587
269. Voas RB, Tippetts AS, Fell JC (2003) Assessing the effectiveness of minimum legal drinking age and zero tolerance laws in the United States. *Accid Anal Prev* 35(4):579–587
270. Wagenaar AC, Holder HD (1991) A change from public to private sale of wine: results from natural experiments in Iowa and West Virginia. *J Stud Alcohol* 52(2):162–173
271. Wagenaar AC, Toomey TL (2002) Effects of minimum drinking age laws: review and analyses of the literature from 1960 to 2000. *Stud Alcohol Suppl* 14:206–225
272. Walden B, McGue M, Lacono WG, Burt SA, Elkins I (2004) Identifying shared environmental contributions to early substance use: the respective roles of peers and parents. *J Abnorm Psychol* 113:440–450
273. Wall TL, Johnson ML, Horn SM, Carr LG, Smith TL, Schuckit MA (1999) Evaluation of the Self-Rating of the Effects of Alcohol form in Asian Americans with aldehyde dehydrogenase polymorphisms. *J Stud Alcohol* 60:784–89
274. Wang JC, Hinrichs AL, Stock H, Budde J, Allen R, Bertelsen S, Kwon KM, Wu W, Dick DM, Rice JP, Jones K, Nurnberger JI, Tischfield JA, Porjesz B,

- Edenberg HJ, Hesselbrock V, Crowe R, Schuckit MA, Begleiter H, Reich T, Goate A, Bierut LJ (2004) Evidence of common and specific genetic effects: association of the muscarinicacetylcholine receptor M2 (CHRM2) gene with alcohol dependence and major depressive syndrome. *Hum Mol Genet* 13:190–1911
275. Wang JC, Hinrichs AL, Stock H, Budde J, Allen R, Bertelsen S, Kwon JM, Wu W, Dick DM, Rice J, Jones K, Nurnberger JI Jr, Tischfield J, Porjesz B, Edenberg HJ, Hesselbrock V, Crowe R, Schuckit M, Begleiter H, Reich T, Goate AM, Bierut LJ (2004) Evidence of common and specific genetic effects: association of the muscarinic acetylcholine receptor M2 (CHRM2) gene with alcohol dependence and major depressive syndrome. *Hum Mol Genet* 13(17):1903–1911
276. Wang PS, Lane M, Olfson M, et al (2005) Twelve-month use of mental health services in the United States: results from the national comorbidity survey replication. *Arch Gen Psychiatry* 62: 629–640
277. Watson AL, Sher KJ (1998) Resolution of alcohol problems without treatment: methodological issues and future directions of natural recovery research. *Clin Psychol: Sci Prac* 5:1–18
278. Weatherburn D, Jones C, Freeman K, Makkai T (2003) Supply control and harm reduction: lessons from the Australian heroin “drought.” *Addiction* 98:83–91
279. Weatherburn D, Lind B (1997) The impact of law enforcement activity on a heroin market. *Addiction* 92(5):557–569, discussion 611–613
280. Wilhelmsen KC, Schuckit M, Smith TL, Lee JV, Segall SK, Feiler HS, Kalmijn J (2003) The search for genes related to a low-level response to alcohol determined by alcohol challenges. *Alcohol Clin Exp Res* 27:1041–1047
281. Williams J, Liccardo Pacula R, Chaloupka FJ, Wechsler H (2004) Alcohol and marijuana use among college students: economic complements or substitutes? *Health Econ* 13(9):825–843
282. Wills TA, Vaccaro D, McNamara G (1994) Novelty seeking, risk taking, and related constructs as predictors of adolescent substance use: an application of Cloninger’s theory. *J Subst Abuse* 6(1): 1–20
283. Wilson RW, Niva G, Nicholson T (1993) Prohibition revisited: county alcohol control consequences. *J Ky Med Assoc* 91(1):9–12
284. Wood E, Tyndall MW, Spittal PM, Li K, Anis AH, Hogg RS, Montaner JS, O’Shaughnessy MV, Schechter MT (2003) Impact of supply-side policies for control of illicit drugs in the face of the AIDS and overdose epidemics: investigation of a massive heroin seizure. *CMAJ* 168:165–169
285. World Health Organization (WHO) (2004) Global status report: alcohol policy. Geneva, Switzerland
286. World Health Organization (1981) Nomenclature and classification of drug- and alcohol-related problems: WHO memorandum. *Bull WHO* 99:225–242
287. Yoon YH, Yi H (2006) Surveillance Report #75: liver cirrhosis mortality in the United States 1970–2003. National Institute on Alcohol Abuse and Alcoholism, Division of Epidemiology and Prevention Research, Bethesda, MD
288. Zimmermann P, Wittchen HU, Höfler M, Pfister H, Kessler RC, Lieb R (2003) Primary anxiety disorders and the development of subsequent alcohol use disorders: a 4-year community study of adolescents and young adults. *Psychol Med* 33: 1211–1222
289. Zuckerman M, Kuhlman D (2000) Personality and risk-taking: common biosocial factors. *J Pers* 68:999–1029