

Contents

List of Figures and Tables Acknowledgements		page vii xi
•	The Journey East	I
	.I Introduction	I
_	.2 The Palaeolithic of Asia	3
Ι	.3 Human Adaptation in the Asian Palaeolithic	5
Chapter 2: 7	he Pleistocene Planet	8
2	.1 Introduction	8
2	.2 The Quaternary Period	9
	.3 Rapid Climate Change	18
2	.4 Conclusion	24
Chapter 3: H	Iominin Dispersal beyond Africa during the Lower	
a	nd Middle Pleistocene	25
3	.1 Introduction	25
3	.2 Lower Pleistocene (2.58–0.775 Ma) Dispersals of Anatomically	
	Archaic Hominins	25
3	.3 Middle Pleistocene (775–127 KBP) Dispersals of Anatomically	
	Archaic Homo	46
3	.4 Dispersal Mechanisms: Faunal Turnovers, Carnivory	
	and Habitat Specificity	62
3	.5 Conclusion	66
Chapter 4: F	Regional Trajectories in Modern Human Behaviour	68
4	.1 Introduction	68
4	.2 The 'Human Revolution' in Europe	68
	.3 Modern Human Behaviour Beyond Europe	80
4	.4 Conclusion	84
Chapter 5: 7	The Initial Upper Pleistocene Dispersal of Homo sapiens	
	Out of Africa	85
5	.I Introduction	85
5	.2 MIS-5e (126–116 KBP)	85
	.3 MIS-5d to the Last Termination (116-22 KBP)	97
5	.4 Conclusion	139

V



Contents

Chapter 6:	Cli	mate, Dispersal and Technological Change during the Last	
-	Ter	mination and Early Holocene in Southeast Asia	142
	6.I	Introduction	142
	6.2	Regional Climate and Conditions across the Last Termination	
		(22–11.7 KBP)	142
	6.3	Genetic Evidence of Human Dispersal during the Last Termination	149
	6.4	Technological Change	150
	6.5	Conclusion	204
Chapter 7:	Tro	pical Subsistence Strategies at the End of the Last Glacial	208
	7.I	Introduction	208
	7.2	Northern Borneo: Niah Caves	210
	7.3	Northern Vietnam: Tràng An	232
	7.4	Eastern Peninsular Malaysia: Gua Sagu and Gua Tenggek	251
	7.5	Inter-Site Patterns of Subsistence through the Last Termination and	
		Early Holocene	261
	7.6	Conclusion	264
Chapter 8:	Ex	Levis Terra	266
_	8.1	Introduction	266
	8.2	The Colonization of New Environments and the Recolonization	
		of Changing Ones	267
	8.3	The Last Termination in Southeast Asia	281
	8.4	Discussion: Hominin Dispersal, Climate Change and Behavioural	
		Evolution	284
	8.5	Conclusion	288
Appendix			291
Bibliography			307
Index			361



List of Figures and Tables

Figures

2.I	Chronostratigraphic correlations for the Quaternary period.	page 11
2.2	Fox Glacier, South Island, New Zealand.	13
2.3	Map showing the locations of the Greenland GRIP, GISP2, NGRIP and	
	NEEM ice-coring operations and the Antarctic locations of the Vostok and	
	EPICA Kohnen Station (EDML) and Dome C ice core drilling sites.	15
2.4	The drill hole at the NEEM ice core drilling camp.	17
2.5	The δ^{18} O GISP2 ice core curve and tropical Atlantic sea surface temperature	
	curve. GIS stages back to c. 57 KBP and GS stages back to c. 40 KBP, with the	
	most recent named stadial and interstadial phases from the European sequence.	19
2.6	An ice floe from the Breiðamerkurjökull glacier, Jökulsárlón lagoon, southeast	
	coast of Iceland.	22
2.7	Schematic of the global thermohaline circulation.	23
3.I	Map showing the East Asian Lower and Middle Pleistocene sites, together	
	with the location of the Chinese Loess Plateau and key stratigraphic reference	
	sections.	27
3.2	Map showing the locations of the Toba and Tambora volcanoes and	
	the distribution of the Australasian tektite strewn field.	43
3.3	Excavations during the 1997 field season at the Gesher Benot Ya'aqov site	
	on the River Jordan, Israel.	47
3.4	Gunung Temporung, Perak, West Malaysia.	57
3.5	Two bifaces from Bukit Bunuh, West Malaysia.	59
4. I	A portrayal of early anatomically and behaviourally modern H. sapiens during	
	our species' Late Pleistocene dispersal through Europe.	69
4.2	Map showing northwest Eurasian sites.	71
4.3	A portrayal of body ornamentation and pigment use (here, on the scalp)	
	among Late Pleistocene H. sapiens around the time of their entry into Europe.	72
4.4	A miniature horse pendant (47 mm in length, 25 mm in height) sculpted	
	from mammoth ivory (perforated between the front legs), dated to	
	the Aurignacian, 30–36 kbp, Vogelherd, Germany.	73
4.5	The panel of horses from the Grotte Chauvet, Ardèche, France.	75
4.6	Franchthi Cave, southeastern Argolid, Greece.	79
4.7	Two N. gibbosulus shell beads from Es-Skhūl, Israel.	81
4.8	A selection of MSA bone points from Blombos Cave, South Africa.	83
5.I	MIS-5 (74-126 KBP) oxygen isotope curve.	87
5.2	Map showing the trans-Sahara watershed, together with key African and	
	Middle Eastern sites.	89
5.3	The Mount Carmel caves viewed from the northwest.	90

vii



List of Figures and Tables

5·4 5·5	The small north-facing Es-Skhūl cave and terrace, Mount Carmel. Map of South, Central and northern Asia, showing the locations of sites	91
	discussed in the text.	95
5.6 5.7	Lang Rongrien rock-shelter under excavation 1984–5. Photograph of the immediate environs of the Bukit Bunuh island site from a	III
	palaeolake, Lenggong Valley, West Malaysia.	113
5.8 5.9	Looking out through the West Mouth entrance to the Niah Caves complex. Looking north across the 'Hell' trench system as preserved today towards	116
	the rock overhang and the Deep Skull still <i>in situ</i> .	117
5.10	A fragment of hominin cranium with pigment adhering to the interior surface.	T T 0
5.11	Pigmented Geoemydidae plastron from Area A of the West Mouth, Niah Caves, dated to c. 41 cal. KBP.	119
5.12	A selection of struck flakes from the Hell and lower E/B1 trenches, Niah Caves.	120
5.13	Photographs of the earliest bone tool from the Hell trench, Niah Caves.	121
5.14	Inter-cutting pit features in the Pleistocene deposits of the West Mouth during the 1959 excavations of this part of the cave and the section sampled	
	by the Niah Caves Project.	123
5.15	Indonesian–Australian excavations at Liang Bua, Flores and deep excavations in Sectors VII and XI.	124
5.16	Excavating LB1 in Sector VII.	125
5.17	Map of Pleistocene Southeast Asia, showing the exposed Sunda Shelf	
	to -116 m, prevailing ocean currents and possible human migration routes through Wallacea.	133
6. I	Map of Pleistocene Southeast Asia, showing the locations of terrestrial and	
	ocean sediment core records.	I44
6.2	Photograph of the Situ Bayongbong location during coring.	145
6.3	Map of Pleistocene Southeast Asia, showing the distribution of bone tool assemblages across the Sunda Shelf, Wallacea and northwest Sahul.	153
6.4	Examples of bone points from the Phase II occupation at Lobang Hangus, classified as likely to have been projectile armatures.	154
6.5	A 'self-barbed' bone point from the Lobang Hangus, Phase II assemblage, compared with a bone-barbed spear from Australia.	155
6.6	An example of a longitudinally split short convex point, or 'gouge', with significant rounding and polishing to the utilised surface, Phase II, Lobang Hangus, and an example of a more complete specimen of this kind of	
6.7	implement from the Late Holocene levels at nearby Kain Hitam Cave. Stingray barb points (probably <i>Pastinachus sephen</i>) excavated by the Niah Caves	156
6.8	Project in Area D of the West Mouth. Evidence of extensively imposed form to create a hafting surface, Ban	157
	Lum Khao.	165
6.9 6.10	Examples of scrapers from the Early Holocene horizons at Lie Siri. Elongated, blade-like scrapers from horizons I and II at Bui Ceri Uato,	169
	East Timor.	170
6.11	Tanged and other retouched blades from Uai Bobo 1, horizons V to IIIc.	171
6.12	Fish hook made from <i>Trochus niloticus</i> , Lene Hara.	172
6.13	Ulu Leang 1 at the end of excavations, July 1969.	178
6.14 6.15	An example of a unifacial, centripally flaked pebble tool or 'Sumatra type',	179
	16×84 mm, from Gunong Pondok, layer G, 3.56 m below datum.	187

viii



List of Figures and Tables

	A pebble tool still <i>in situ</i> in the Deep Sounding of the Hang Boi excavation.	189
6.17	Limestone blade-flake and core face rejuvenation flake from Hang Trông,	
6 - 0	Tràng An, Vietnam. An early stage in excavations at Gua Kerbau in 1926.	191
6.18	Photograph showing excavation work beneath the rock overhang (Area A,	195
7.I	Niah Caves Project), probably 1965.	211
7.2	The archaeological compound of the West Mouth, Niah Caves.	211
7.2	Fragments of Cercopithecidae long bone constituting manufacturing debris	221
7.3	from bone tool production.	227
7.4	Examples of cut-marked distal humeri from Cercopithecidae from the	227
7.4	*	220
	Lobang Hangus, Phase II occupation. The NISP values and frequency of occurrence of butchery traces to the distal	229
7.5	<u> </u>	
	humerus of Arctictis binturong compared to other key taxonomic groups from	227
- 6	the Phase II cut-marked assemblage from Lobang Hangus.	231
7.6	A view of the Upper Cave at Hang Boi looking east towards the	
	archaeological compound and access to the Lower Cave, with the opening to the cave mouth visible.	222
		232
7.7	Contexts and sample columns in the east and south-facing sections of Trench	222
- 0	I at Hang Boi. The southeast-facing section of excavations at Hang Boi (2009 field season),	233
7.8		
	illustrating the whole shell and sediment-rich crushed shell stratigraphy that	
	has been revealed in the Upper Cave shell midden.	235
7.9	The remains of a hunter's small fire found on the open floor of the Upper	220
	Cave at Hang Boi in 2010.	238
7.10	Three pierced Neritidae shells recovered from midden contexts at Hang Boi.	239
7.11	Villopotamon sp. Raffles Museum of Biodiversity Research at the University of	2.40
	Singapore and specimens of archaeological chelae from Hang Boi.	240
7.12	Panthera cf. pardus specimen from Hang Boi compared against Panthera tigris.	241
7.13	Archaeological (L) maxilla fragment with PM ¹ , M ¹ and M ³ in situ, compared	
	alongside Ratufa (= Sciurus) bicolor.	242
7.14	Keeled box turtle (Cuora mouhotii) and a nuchal fragment identified to this	
	species.	245
7.15	One of a population of <i>Macaca arctoides</i> (stump-tailed macaque) that currently	
	inhabits Tràng An Park.	247
7.16	Asian leaf turtle (Cyclemys dentata) and specimens of third and fourth (L)	
	peripheral dermal plates of this species from Gua Sagu.	257
7.17	Cut-marks to the inside of the hyo- or hypoplastron dermal plate of a	
	chelonid, Gua Sagu, and cut-marks to a chelonid femur or humerus	
0	mid-shaft.	259
8.1	View of wet rice fields just prior to the transplantation of rice seedlings, near	
	the village of Pa' Dalih in the Kelabit Highlands of Sarawak.	275
_		
Tab	les	
3.I	Identified Mammalian Fauna from Goudi/Majuangou III and	
J. 1	Xiaochangliang, Northeast China	33
3.2	Identified Mammalian Fauna from Chenjiawo and Gongwangling,	33
J.2	Central China	36
3.3	Identified Mammalian Fauna from the M3 and M4 Components of	50
J·3	the 'Yuanmou Hominid Fauna', Yuanmou Basin, North Central China	38
	The second secon	50
		•
		ix



List of Figures and Tables

7. I	Identified Taxa from the Early and Late Phase II Occupation of Area A, in	
	the West Mouth of the Niah Caves (Excluding Human Remains)	213
7.2	Taxa Represented during the Early Holocene Occupation of the West	
	Mouth, Area D	222
7.3	Taxa Represented during the GIS-1 Occupation of Lobang Hangus	224
7.4	Identified Invertebrate Macrofaunal Remains from the Upper Cave Shell	
	Midden (2007–8 Samples), Hang Boi	237
7.5	Vertebrate Fauna Identified and Quantified from the Cave Mouth Midden	
	at Hang Boi (2008–9 samples quantified)	243
7.6	Identified Vertebrate Fauna from Pleistocene and Early Holocene Levels	
	at Gua Sagu (spits 5–11) and Gua Tenggek (spits 4–7)	255
8.1	The Broad Adaptive Circumstances and Strategies of Tropical	
	Hunter-Gatherer Communities during most of the Last Termination,	
	Southeast Asia	283