
Contents

<i>Preface</i>	<i>page</i> vii
<i>Acknowledgements</i>	ix
<i>Introduction</i>	1
1 The occurrence of dormancy in the Gramineae	3
1.1 Wild grasses	6
1.2 Forage grasses	7
1.3 Cereal grasses	19
1.4 Weed grasses	30
1.5 Dormancy and persistence of individuals and populations	34
1.6 Xerophytes, mesophytes and hydrophytes	42
1.7 Terminology and definitions of dormancy	43
1.8 A conceptual approach for understanding seed dormancy	48
1.9 <i>Avena fatua</i> , a prototype model of grass seed dormancy	59
2 Mutual influences of inflorescence and caryopsis parts on dormancy	60
2.1 Position of the caryopsis on the parent plant	60
2.2 Lemma, palea, glumes and awns	65
2.3 The grain coat – pericarp and testa	83
2.4 Embryo and scutellum	94
2.5 Interrelation between embryo and maternal storage tissue	106
3 Environmental influences on seed dormancy	114
3.1 Water – liquid and vapour phases	114
3.2 Effects of radiation – heat/cold, light, microwaves, magnetism, photoperiodism	119
3.3 Latitudinal influences – tropical and temperate zones	158
3.4 Gases – oxygen and anoxia, carbon dioxide, ethylene, volatile organics; pressure	161
3.5 Growth regulators – promoters, inhibitors, limiting and allelopathic factors	170
3.6 Soil and agricultural practice	187

<i>Contents</i>	
4	Timing of dormancy 195
4.1	Semantic considerations 195
4.2	Timing in the induction, maintenance and release of dormancy 197
5	Modelling the induction, maintenance and termination of dormancy in grass seeds 207
5.1	Modelling theory for biological systems 207
5.2	A ‘black box’ perspective of seed behaviour 210
5.3	An energy flow model for grass seed dormancy 218
5.4	Similarity and diversity among dormancy states in grass seeds 224
5.5	Conclusion 228
	<i>Bibliography</i> 232
	<i>Index</i> 281